
SOLUTIONS OF NONLINEAR MATRIX EQUATIONS(U) AIR FORCE INST OF TECH WRIGHT-PATTERSON AFB OH SCHOOL OF ENGINEERING B W COLLETTI DEC 85 AFIT/GOR/MA/85D-1 F/G 12/1 AD-A163 841 1/2 UNCLASSIFIED NL



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A



SOLUTIONS OF

NONLINEAR MATRIX EQUATIONS

Thesis

Bruce W. Colletti Captain, USAF

AFIT/GOR/MA/85D-1

DISTRIBUTION STATEMENT A

Approved for public releases

Distribution Unlimited

DEPARTMENT OF THE AIR FORCE

AIR UNIVERSITY

AIR FORCE INSTITUTE OF TECHNOLOGY

Wright-Patterson Air Force Base, Ohio

FEB 1 0 1986



AFIT/GOR/MA/85D-1



SOLUTIONS OF

NONLINEAR MATRIX EQUATIONS

Thesis

Bruce W. Colletti Captain, USAF

AFIT/GOR/MA/85D-1

Approved for public release; distribution unlimited

SOLUTIONS OF NONLINEAR MATRIX EQUATIONS

Thesis

Presented to the Faculty of the School of Engineering
of the Air Force Institute of Technology
Air University
in Partial Fulfillment of the
Requirements for the Degree of
Master of Science



bу

Bruce W. Colletti, B.S.

Captain, USAF

December 1985

Acces	ion For	7	
DTIC	lounced		
By Distribution /			
Availability Codes			
Dist	Avail alid Specia	/ or	
A-1			

Approved for public release; distribution unlimited

Preface

My earliest memory of mathematics recalls writing numbers in the spaces of a grid. Little did the first grader know that by creating this "box of numbers", he was laying the groundwork for subsequent study in matrices, themselves very special "boxes of numbers." After studying matrix theory as a high school freshman, I remember hoping for a chance someday to pursue it in great depth. Though the hope has now been realized, it will by no means be put to rest: for my master's in mathematics, I will pick up where this thesis left off.

Fortune doesn't always favor us, and as Machiavelli says, will strike after being wooed and cajoled. However, she must have been in a good mood in the Fall of '84 because our class had Dr. John Jones Jr for Numerical Analysis. His presentation of matrix algebraic concepts delighted me, and upon discovering his abiding interest and renown with matrix equations, decided to study under him. Undoubtedly, Dr Jones was one of two teachers having a profound impact on my mathematical development. I couldn't have wished for a finer advisor and coach than him. His patience and positive outlook sustained me during those times when I was setting back mathematical science instead of advancing it. For him, theory and applications are one, though primacy goes to theory. Just as it should be. Thanks for your interest in me, Dr Jones. I will return your concern years hence to one

of my students.

To LTC (and Dr.) Charles Ebeling, my reader in the Operations Research Department, I want to thank for being (I believe) a mathematician at heart, and hence an ally in the department. If I couldn't have done a mathematical thesis, I would have done a simulation one and sought you out as an advisor. Thanks for not discouraging me from pursuing an interest begun years ago.

Contents

	Pag e	
Preface	ii	
List of Figures	v	
List of Symbols	٧i	
Abstract	vii	
I. Overview and Literature Review	1.1	
II. Foundations	2. i	
Lypanov Equation	2.2 2.3 2.5 2.11	
III. Solving the Linear Equation AX = b	3.1	
AX = b, rank(A) constant	3.4 3.12 3.16	
IV. Solutions to Lypanov and Second Order Riccati Equations	4.1	
Lypanov Equation	4.1	
Tensor Solution	4.1	
Transformation	4.5	
Second Order Riccati Equation	4.16	
V. Solutions to Third Order Riccati Equation	5.1	
VI. Conclusions and Recommendations	6.1	
Appendix A: BASIC program performing matrix operations	A. 1	
Bibliography		
Vita	BIB-4	

<u>List of Figures</u>

Figure		Page
1.	Network tree representing the polynomial	
	$3x^2yz^{13} - 7x^{12}y^7z + 8x^{12}y^7z^{22}$	A.8
2.	Branch for x²	A.9
3.	Branch for x2 + x2y	A.9
4.	Branch for $x^2 + 2x^2y$	A.9
5.	Branch for $x^2 + 2x^2y + y^2$	A.9
6.	Session building matrix (A.10)	A. 10
7.	Session viewing the matrix fA(R) preceeding	
	(4.23)	A. 11

<u>List of Symbols</u>

Omxn	an m X n matrix whose entries are all zero
A A B	tensor product of matrices A and B
Amxn	the m X n matrix A
¢	the set of complex numbers
¢(P)	the set of polynomials with coefficients in the complex numbers and variables in the components of the n-tuple $\ensuremath{\Gamma}$
¢ m×n(⊣)	the set of m X n matrices whose entries are polynomials with coefficients in the complex numbers and variables in the components of the n-tuple $\ \ \cap$
¢(x)	the set of polynomials in the variable x
¢(x,y)	the set of polynomials in the variables x,y
det (A)	the determinant of the matrix A
iff	if and only if
Inxn	the n X n identity matrix
rank(A)	the rank of the matrix A
S ₁ - S ₂	the set of elements which are in the set $S_{\text{\tiny 1}}$ but not in the set $S_{\text{\tiny 2}}$
Г	an arbitrary n-tuple (רוּ,,רח)
× (S	the entity x is contained in the set S
Z+	the set of positive integers 2 0

<u>Abstract</u>

This paper seeks the solutions to a system of equations (equalities) in n variables by expressing the system in matrix algebraic form. Properties of the solutions to the ensuing matrix equation are investigated using similarity transformations. The three types of matrix equations to be studied are: the linear equation.

AX = b

the Lypanov equation

$$AX - XB = C$$

the second-order Riccati equation -

$$XDX + AX + XB + C = O_{j}^{-1}$$

and the third-order Riccati equation -

$$XAXBX + XCX + DX + XE + F = 0$$

The entries of all matrices, including the solution X, are restricted to being polynomials in Γ having complex coefficients, where Γ is the n-tuple of indeterminates. That is, all matrices are elements of the ring $\mathfrak{t}^{\mathsf{mxn}}(\Gamma)$ for m and n of appropriate size.

Because adding and multiplying matrices (having multivariate polynomial entries) is tedious in practice, an interactive BASIC program is presented in the appendix. This program, which can be used on a personal computer, permits the user to perform operations on matrices having multivariate polynomial entries. Via menu selections, the user may perform

-weighted addition between two matrices

- -multiplication between two matrices
- -create matrices, with an option of building a diagonal matrix whose diagonal entries are all equal
- -view matrices
- -transpose a matrix
- -extract special submatrices (U,M,V,N of Chapter IV) from a given matrix.

I Overview and Literature Review

Overview. This paper seeks the solutions of the third-order Riccati matrix equation

$$XAXBX + XCX + DX + XE + F = 0$$
 (1.1)

where the entries of the matrices A,B,C,D,E,F,X are multivariate polynomials. In solving (1.1) the linear matrix equation

$$AX = b$$

the Lypanov matrix equation

$$AX - XB = C$$

and the second-order Riccati matrix equation

$$XDX + AX + XB + C = O$$

will be addressed because the form of their solutions will hint at the nature of those to (1.1). Because cursory knowledge of matrix algebra is sufficient to motivate the paper's thesis—the solution to (1.1)—the rigorous (and lengthy) definition of terms and statement of objective will be made in the next chapter.

Why a matrix equation (such as those given above) is worthy of attention, let alone finding its solution, is a legitimate concern. Practical problems often arise which require the solution to a system of m equations in n unknowns, e.g., the system of equations

$$ax + by = c$$
 (1.2)
 $dx + ey = f$

where all variables except x and y are known. One way to

find the solutions \mathbf{x} and \mathbf{y} is to express the system in its equivalent matrix form

$$\begin{bmatrix} a & b \\ d & e \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} c \\ f \end{bmatrix}$$

(which is a linear matrix equation) and solve for x and y, i.e.

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a & b \\ d & e \end{bmatrix} \begin{bmatrix} c \\ f \end{bmatrix}$$

Thus, a system of equations may give rise to a matrix equation whose solution in turn gives the answer to the original system.

Other more complex systems may have a matrix representation. For instance

$$\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & b \\ c & d_{3} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} e \end{bmatrix}$$

$$= \begin{bmatrix} xa + yc & xb + yd \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$= \begin{bmatrix} x(xa + yc) + y(xb + yd) \end{bmatrix}$$

$$= \begin{bmatrix} x^{2}a + xy(c + b) + y^{2}d \end{bmatrix}$$

which means that the system consisting of the one equation

$$x2a + xy(c + b) + y2d = e$$

is represented by the matrix equation (1.3). These two examples show that matrix equations warrant attention if for no other reason than they can help solve systems of equations arising in practical problems. Indeed, given the nonlinear optimization challenge

Maximize f(x) subject to

(1.4)

equality constraint,

equality constraint,

the constraint set may be expressable as a matrix equation. If so, this equation can be solved to identify the feasible region determined by the constraints and thence the optima. This approach is an alternative to solving (1.4) using, say, Lagrangian multipliers and partial differential calculus.

In examples (1.2) and (1.4), the variables are often taken to represent real or complex numbers. However, there is nothing stopping the variables from assuming functional values. For instance (1.2) may assume the form

$$a(u,v)x + b(u,v)y = c(u,v)$$
 (1.5)
 $d(u,v)x + e(u,v)y = f(u,v)$

where a,b,c,d,e,f are functions of the parameters u and v. Though the solutions x and y of (1.5) wouldn't necessarily have numeric values, this is all right because all other variables are functions: x and y will likely take on functional forms. To illustrate,

$$\begin{bmatrix} u^{2}v & u \\ uv & v \end{bmatrix} \begin{bmatrix} u + v \\ u - v \end{bmatrix} = \begin{bmatrix} u^{2}v + u^{2}v^{2} + u^{2} - uv \\ u^{2}v + uv^{2} + uv - v^{2} \end{bmatrix}$$

and so a solution to the matrix equation

$$\begin{bmatrix} u^{2}v & u \\ uv & v \end{bmatrix} \begin{bmatrix} x(u,v) \\ y(u,v) \end{bmatrix} = \begin{bmatrix} u^{3}v + u^{2}v^{2} + u^{2} - uv \\ u^{2}v + uv^{2} + uv - v^{2} \end{bmatrix}$$

is x(u,v) = u + v and y(u,v) = u - v. As given in (1.1), this paper will address matrix equations whose entries

assume the functional form of a polynomial in several variables.

Another area where matrix equations appear (where the matrices' entries are multivariate functions) is control theory, the science addressing the orbital natures of objects in space. Matrix equations frequently arise from systems of differential equations which describe a satellite's orbit. For example, given X(t) a columnar matrix whose entries (each describing an aspect of a satellite's orbit) are functions in the parameter t, a theorem from control theory states that the equilibrium of the system of differential equations

is asymptotically stable if the Lypanov matrix equation

$$F^{T}P + PF = -C \qquad (1.6)$$

has a positive-definite solution P for any matrix $\mathbb{C} > 0$. Though the terms in this theorem won't be defined here (see [4:144-152]), the important item is that a satellite's orbital stability depends on the solution of a Lypanov matrix equation (1.6).

Matrix equations can also describe the path taken by an X-ray passing through matter. This is a concern of the medical community since the quality of CAT scan pictures depends on the manner in which the X-rays penetrate the skull. A good description of the path may allow finding a

brain tumor, whereas a poorly chosen angle of entry may not.

R. Vasudevan [32] develops higher order Riccati equations describing how beam particles scatter upon hitting matter, e.g. the skull. Simply put, scattering is modelled by including higher powers of a solution matrix X in a Riccati equation. For instance, scattering may be roughly described by the second order Riccati equation

\$ P.

0.

$$XDX + AX + XB + C = 0$$

and more completely by the third order Riccati equation

$$XAXBX + XCX + DX + XE + F = O$$

where the coefficient matrices A,B,C,D,E,F reflect known characteristics of the environment in which the X-rays behave. Higher powers of X will more accurately describe scattering. Bellman and Vasudevan [2] describe techniques reducing a given Riccati equation to one of lower form. This reduction (quasi-linearization) leads to a series which converges to the actual solution of the original equation.

<u>Literature Review</u>. The literature supporting this paper addresses three topics:

- I generalized inverses of matrices
- II algebraic structures of matrices over fields and rings
- III the Lypanov and Riccati equations

Category I. An understanding of matrix generalized inverses launched the research behind this paper. Matrix generalized inverses encompass the traditional notion of an inverse by assigning these to non-square matrices. The theory is well developed for

matrices whose entries are complex numbers, and is described below.

The physicist R. Penrose [28] proved in 1955 that for any matrix A whose entries are complex numbers, there exists matrices U,Y,Z,W having complex entries such that:

- $1. \quad AUA = A \tag{1.7}$
- 2. YAY = Y
- AZ equals its conjugate transpose (i.e., AZ is Hermitian)
- 4. WA is Hermitian

The matrices U,Y,Z,W are known as the "generalized inverses" of A. Penrose also showed the existence of a matrix X which simultaneously satisfied conditions (1) thru (3), and a Y simultaneously satisfying (1), (2) and (4). Penrose also showed that if a matrix X satisfied all conditions (given a matrix A), then X was unique. From the traditional matrix algebra viewpoint, this unique X corresponds to the familiar inverse of a square matrix A whose determinant is not equal to zero. Though Penrose's work was original, he was unaware that the mathematician E.H. Moore [24] had proved (using abstract algebra) the existence of generalized inverses for arbitrary rings more than three decades earlier.

Rao and Mitra's textbook [29] is devoted to the study of generalized inverses, and is a frequently cited authority in this area. The first two chapters of [6] present techniques (with the justifying theorems) generating various types of generalized inverses for a given matrix A, while Captain Craig Murray, AFIT Class GCS 85D, has recently

completed the programming of these techniques. Two papers, [12] and [21], use generalized inverses to prove theorems about special types of matrix equations whose entries are complex numbers.

The power of generalized inverses weakens as one moves from matrices with complex entries to those with multivariate entries, the subject of this paper. This is because the theory in the latter is still very young. Sontag [31] discusses the existence of some generalized inverses for special matrices, while Jones [11] uses these generalized inverses to solve certain matrix equations.

Category II. Most of this paper's research dealt with the decisive influence on the existence and form of solutions to matrix equations: the algebraic structure of matrices. An illustration of the importance of algebraic structure is readily given: the solution of the equation

$$x + 7 = 5$$
 (1.8)

cannot be found among the set of positive integers Z^+ (even though the coefficients 7 and 5 are positive integers) because its structure doesn't include negative numbers. Since -2 is the solution, (1.8) must be placed in a larger algebraic structure in order for a solution to exist. Thus one moves from Z^+ to the set of all integers Z^- . Though Z^- has a larger algebraic structure than Z^+ it still isn't large enough to contain the solution of the equation

$$2x - 8 = 7$$
 (1.9)

even though the coefficients 2,-8,7 are found in Z. It now becomes necessary to move to a larger structure (the set of rational numbers) to find the solution to (1.9), 15/2. Likewise, to solve the equations $x^2 = 2$ and $x^2 + 1 = 0$ one must move to the larger algebraic structures found in the irrational and complex numbers respectively.

Matrices too lend themselves to different structures. However, not only must the form of a matrix be addressed. (e.g., whether it's invertible, whether it's similar to another matrix), but the structure in which individual entries are found must also be considered. For instance, decomposing a matrix into a product of other matrices may depend upon whether or not an entry is irreducible in its own setting. A trivial example is decomposing the 1 X 1 matrix

$$[x^2 + x - 1] \tag{1.10}$$

into a product of two other 1 X 1 matrices each having polynomial entries whose coefficients are integer, as is the entry in (1.10). Such a decomposition is impossible because $x^2 + x - 1$ cannot be expressed as a product of two polynomials with integer coefficients. Indeed, the roots of this equation are $(-1 \pm 45) / 2$, which are irrational.

In general, this paper allows matrix entries to be multivariate polynomials (having complex coefficients, e.g., $x^2 + xy$, $xy^2z - (2 + 3i)x^6yz$), who in turn have complex algebraic structures. In fact, a solid understanding of

abstract algebra would have helped make several articles intelligible. Though Fraleigh's text [7] was found to be an excellent primer on abstract algebra, time did not allow sufficient study of a subject key to this paper. Wang [33] gives an algorithm for irreducible factoring of multivariate polynomials having coefficients in an arbitrary algebraic number field. McClellan [23] presents methods for solving systems of equations involving univariate polynomials with rational coefficients (this paper highlights his doctoral dissertation). Two papers from the early twentieth century, [3] and [27], discuss the form of factors between polynomials.

The algebraic structures of matrices was addressed by Frost and Storey [8] and Lee and Zak [20]. These special structures, called Smith Forms, are discussed in the next chapter. Unfortunately, only the Smith Forms of matrices having bivariate polynomial entries can be addressed: the Smith Form involving multivariate entries remains an unresolved issue in mathematics.

Category III. Working from a base rooted in generalized inverses and recognizing the critical role of algebraic structures on matrices (and again among their entries), a study of matrix equations can proceed. Though much has been published on finding solutions to systems of equations, many authors succeed in finding only particular solutions. After all, finding the general forms of all

solutions may have either been a far too ponderous task or the mathematical approach proved elusive. In any case, viewing a system of multivariate equations as a single matrix equation (having a solution in its own right) is not commonplace. Depending on the nature of the system, the entire solution set may be found by solving a matrix equation representing the system.

Work has been done on matrix equations having complex entries. Roth [30] (a frequently cited paper) wrote on the Lypanov matrix equation in 1952, and had his work extended in 1972 by Jones [12] who also addressed the second order Riccati equation, using generalized inverses to identify solutions. Morris and Odell [25] attempted to find the common solutions to a set of linear matrix equations $\{A_kX = B_k\}$ by using generalized inverses. Lancaster [18] provides several approaches (none using generalized inverses) to solving the matrix equation

As matrix equations assume multivariate polynomial entries, the constraints cited in the previous section seriously handicap the search for solutions. As a result, pioneering work in these equations is still underway.

The papers [11], [16] and [17] give recent results connected with the Lypanov and second/third order Riccati equations.

Collectively, these papers address the roles of generalized inverses and matrix algebraic structures in the search for

solutions to these matrix equations.

II Foundations

This chapter lays the groundwork for later developments. Because this paper seeks solutions to matrix equations, the latter's nature will be addressed first.

Matrix Equations. A matrix equation is an algebraic expression whose arguments involve matrices. Matrix equations are frequently used to represent a kindred system of equations. For instance, the system of equations

$$2x + 3y = 7$$

 $4x - 2y = 1$

is represented by the matrix equation

$$\begin{bmatrix} 2 & 3 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 \\ 1 \end{bmatrix}$$

which has the familiar form

$$AX = b \tag{2.1}$$

where

$$A = \begin{bmatrix} 2 & 3 \\ 4 & -2 \end{bmatrix} \qquad X = \begin{bmatrix} x \\ y \end{bmatrix} \qquad b = \begin{bmatrix} 7 \\ 1 \end{bmatrix}$$

The reason for expressing a system of equations in matrix algrebraic form is, of course, so that the machinery of matrix algebra can be used to find a solution to the given initial system.

Other systems of equations have matrix representations. For instance, the system

$$x + 2u = 5$$
 $3x + 4u = 7$
 $9x + 10u = 23$
 $y + 2v = 6$
 $3y + 4v = 8$
 $9y + 10v = 24$
 2.1

has the matrix form

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 9 & 10 \end{bmatrix} \begin{bmatrix} x & y \\ u & v \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 7 & 8 \\ 23 & 24 \end{bmatrix}$$

which in turn has the form (2.1) where

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 9 & 10 \end{bmatrix} \qquad X = \begin{bmatrix} x & y \\ u & v \end{bmatrix} \qquad b = \begin{bmatrix} 5 & 6 \\ 7 & 8 \\ 23 & 24 \end{bmatrix}$$

Thus in viewing (2.1), it isn't necessary that b and X be one-columnar matrices and A square, as is traditionally presented.

Lypanov Equation. The linear matrix equation (2.1) leads to the slightly more complex equation

$$AX + XB = C (2.2)$$

which is known as the Lypanov equation. In order for (2.2) to be defined, the matrices A,B must be square (though not necessarily of the same size) and the matrices X,C must share the same size (though they may be rectangular). To see why this is true, let the following hold:

For AX to be defined, b must equal c: b = c. For XB to be defined, d = e. AX is a X d and XB is c X f. For AX + XB to be defined, a = c and d = f. Since b = c and a = c, a = b and A is square c X c. Similarly for B: d = e and d = f and so e = f which makes B square d X d. It follows that AX + XB is c X d which is the size of X.

To illustrate (2.2), the system

$$(x + 2u) + (5x + 8y + 11z) = 14$$
 (2.3)
 $(y + 2v) + (6x + 9y + 12z) = 15$
 $(z + 2w) + (7x + 10y + 13z) = 16$
 $(3x + 4u) + (5u + 8v + 11w) = 17$
 $(3y + 4v) + (6u + 9v + 12w) = 18$
 $(3z + 4w) + (7u + 10v + 13w) = 19$

has the Lypanov form

$$\begin{bmatrix}
1 & 2 \\
3 & 4
\end{bmatrix}
\begin{bmatrix}
x & y & z \\
u & v & w
\end{bmatrix}
+
\begin{bmatrix}
x & y & z \\
u & v & w
\end{bmatrix}
\begin{bmatrix}
5 & 6 & 7 \\
8 & 9 & 10 \\
11 & 12 & 13
\end{bmatrix}$$

$$= \begin{bmatrix}
14 & 15 & 16 \\
17 & 18 & 19
\end{bmatrix}$$

Of course, (2.3) can be expressed as (2.1) by combining like terms, thereby creating six equations in six unknowns.

However, it will be seen (Chapter IV) that by solving the Lypanov equation instead of its equivalent linear form (2.1), solutions of a higher order matrix equation can be derived, one in which the Lypanov is embedded. This higher equation is known as the second-order Riccati equation.

Riccati Equation and Higher. In high school algebra, a student moves from solving the simple linear equation bx = c to the higher order quadratic equation $ax^2 + bx = c$ (in which is embedded the linear equation bx = c). In the present setting, the former equation is likened to the Lypanov matrix equation and the second-order Riccati matrix equation to the quadratic. The Riccati equation has the form

$$XDX + AX + XB + C = 0 (2.4)$$

For (2.4) to be defined, A and B must be square (possibly

not the same size), D must have the size of X^T , and X, C share the same size. The argument for this follows that of the Lypanov equation.

To illustrate (2.4), the Riccati form of the system

$$x^{2} + uy + 2x = -1$$
 (2.5)
 $xy + yv + 2y = -2$
 $xz + yw + 2z = -3$
 $ux + vu + 2u = -4$
 $uy + v^{2} + 2v = -5$
 $uz + vw + 2w = -6$

i s

$$\begin{bmatrix} X & Y & Z \\ U & V & W \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} X & Y & Z \\ U & V & W \end{bmatrix} & + & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} X & Y & Z \\ U & V & W \end{bmatrix} & + & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} X & Y & Z \\ U & V & W \end{bmatrix} & + & \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} & + & \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} & = & \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

As with the Lypanov equation, there may exist many solutions to the second-order Riccati equation. In fact it will be seen in Chapters III and IV that the solutions of (2.4) will satisfy pairs of equations determined by the matrices A,B,C,D in (2.4).

As it is an easy matter to create a polynomial of a given degree (e.g., $3x^5 + 3x^2 - 7$), so is it also to create a higher order matrix equation. For instance, the third order Riccati equation

$$XAXBX + XDX + AX + XB + C = 0$$
 (2.6)

or the related matrix equation

EXAXBXF + GXDXH + AXJ + KXB +
$$C = 0$$

where the matrices not equal to X are known and so can be 2.4

) •

thought of as coefficients of a polynomial in x. However, the non-commutativity of matrix multiplication gives the placement of each coefficient matrix a decisive influence on the solution.

A natural question that arises is whether or not a given system has an accompanying matrix form. At present, the only known way to answer this question is to experiment, an understandably unattractive task. However, if a matrix representation is derived (or stumbled upon), the concepts presented in this paper will help to identify the solutions to the given matrix equation, and thus to its underlying system of equations.

Matrices Over a Ring. Two items determine the nature of a matrix: its size and entries. Concerning the latter, a matrix is said to be "over a set S" if and only if its entries come from the set S. For example if $S = \{1,2,3\}$ then the 1 X 4 matrix

$$A = [3 1 1 2]$$

is "over S" because each of its entries belong to S. However, the 2 X 3 matrix

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 2 \end{bmatrix}$$
 (2.7)

is not "over S" because B(2,3) = 0, which is not a member of S. If however, $S = \{\text{the real numbers}\}$, then B is "over S". Likewise, if

$$A = \begin{bmatrix} 2 + 3i & 7 \\ 9 & 8 - 7i \end{bmatrix}$$
 (2.8)

then A is not a matrix over the real numbers, but is a matrix over the complex numbers. In this paper, the set of complex numbers will be given by the symbol ¢ (see the "List of Symbols").

These examples lead to a notation used throughout this paper: given a set S, the set of all m X n matrices whose entries are in S is symbolically given by

If A is an m X n matrix whose entries are in S, then the statement "A is an element of $S^{m\times n}$ " is represented by

A & Smxn

Example: From (2.8), A $\notin \mathfrak{c}^{2\times 2}$ because A is a 2 X 2 matrix whose entries are complex. However from (2.7), B is not in $\mathfrak{c}^{2\times 2}$ because B is a 2 X 3, despite the fact that the entries of B are elements of the complex numbers (the set of real numbers is a subset of the complex numbers). However, B $\notin \mathfrak{c}^{2\times 3}$

The concept of "overness" will now be extended. Let

Thus if $S = \{1, 2, 3\}$, then

$$f(x) = x^2 + 2x + 3 \in S(x)$$

because f(x) is a polynomial in x with coefficients in S. However

$$f(y) = y^2 + 2y + 3$$

is not in S(x) because f(y) is a polynomial in y. Likewise, $f(x) = 7x^2 + 2x + 3$

is not in S(x) because the coefficient of x^2 is seven, which is not found in $S = \{1,2,3\}$. It follows that $\xi(x)$ is the set of polynomials (in the variable x) with complex coefficients.

Similar to (2.10), let

S(x,y) = {polynomials in the variables x,y whose coefficients are in the set S}

Thus if $S = \{4,5,2\}$, then

 $f(x,y) = 4x^{10}y^{5} + 2x^{2}y + 5x + 5y + 5(x,y)$ because f(x,y) is a polynomial in x,y with coefficients in S. It follows that f(x,y) is the set of polynomials in the variables x,y with complex coefficients.

In general, let Γ represent the n-tuple (x_1,x_2,\ldots,x_n) . Then given a set S, $S(\Gamma)$ is defined to be the set of all polynomials (in the variables found in Γ) with coefficients in the set S.

Example: If $\Gamma = (x,y,z,v,w)$, then $\mathfrak{C}(\Gamma)$ is the set of all polynomials in the variables x,y,z,v,w with complex coefficients. Thus

 $f(x,y,z,v,w) = (4+5i)x^2 + xyzvw - (2+12i)w^7 + f(r)$. However if S is the set of real numbers, then f(x,y,z,v,w) is not in S(r).

Now to extend notation to matrices. Define

 $S^{m \times n}(r) = \{m \times n \text{ matrices whose } (2.11)$ entries are in $S(r)\}$ Thus if $S = \{1,2,4,8,9\}$ and r = (x,y,z) and

$$A = \begin{bmatrix} x^2yz + 9z & x \\ 2x & 4yz + 8z^2 \end{bmatrix}$$

then A f $S^{2\times 2}(\Gamma)$ because A is 2 X 2, and each entry of A is a polynomial in $\Gamma = (x,y,z)$ with coefficients in S. However if $S = \{1,2,4,8\}$, then A would no longer be in $S^{2\times2}(r)$ because a coefficient of A(1,1) isn't in S (namely, 9). From (2.11), $\mathfrak{q}^{m\times n}(n)$ is the set of all m X n matrices whose entries are polynomials in r having complex coefficients.

With this final definition, the object of this paper can be stated: find the solution $X \in C^{m \times n}(\Gamma)$ of the third-order Riccati equation

XAXBX + XCX + DX + XE + F = 0(2.12)given of the tuple of indeterminates in the polynomial entries of A,B,C,D,E,F ($\phi^{k\times j}(\cap)$ for k,j,m,n of appropriate size.

One may feel that in (2.12), mandating X & &(r) unduly restrictive. After all, forcing the solutions of $x^2 - 2 = 0$ (a polynomial in x over the integers, i.e., a polynomial with integer coefficients) to again be a polynomial over the integers isn't possible, since 12 is irrational. Also, given the matrix equation AX = b with the entries of A,b integer and A square and invertible, it may not be possible to have the solution $X = A^{-1}b$ to again have integer entries. Why then require X in (2.12) to have entries in ¢(n) when (like these two analogies) it may be

necessary to leave ¢(n)? The argument is well posed, since a given matrix equation may indeed not have a solution in a given space. In fact, practical solutions could be missed by restricting solutions for theoretical reasons.

The reasons for mandating X (cmxn(r) are in part pragmatic. First of all, polynomial entries don't lend themselves to singularities: X would be defined at all values of r. This is important because given an entry that is a rational function (e.g., 1/x), there may exist a singularity or discontinuity at a value which in fact does have physical significance. Because of the rational entry, the form of the solution may prove untenable. Secondly, many functions can be approximated by a series of polynomials, e.g., Chebysev polynomials [5:239]. In fact, computational considerations may force the analyst to use polynomial approximations of a function. Restricting the entries of X to ¢(r) would help generate these polynomials to be used in approximating the solutions to a phenomena.

The driving force, however, for requiring X $\notin \mathfrak{t}^{m \times n}(\Gamma)$ is theoretical: is it possible to find some solutions in the same space as the coefficient matrices in a given matrix equation? If so, how would the solutions be obtained? It may very well be that solutions thus found may shed light on the nature of solutions which lie outside the required space.

A word needs to be said on the setting for addition and

multiplication on matrices over $\mathfrak{C}(r)$. Though these operations seem to hardly merit concern, there is in fact an extremely important algebraic structure behind them. Some of the current literature on matrix equations refers to this "ring" structure of $\mathfrak{C}^{m\times n}(r)$.

In the broad setting, a set of objects S (e.g., matrices) is given, and two operations, + and *, are defined (e.g., matrix addition and multiplication) between a pair of elements of S. These operations have the property that if a,b (S then: a + b , a * b (S and a + b , a * b assume unique values (e.g., a + b has one value and one value only). Operations with these closure and uniqueness properties are called binary operations. Depending on the nature of the operations, an algebraic structure (denoted by (S,+,*) is defined on the set S taken together with the operations. For S the set of matrices over ¢(n) and the customary operations of matrix addition and subtraction, (S,+,*) is known as a ring.

A ring $\langle R,+,* \rangle$ is a set R together with two binary operations + and * defined on R such that the following axioms are satisfied [7:195]:

It is an easy matter to confirm that the operations of matrix addition and multiplication satisfy the ring axioms. 2.10

11

Rank and Determinant of a Matrix. This chapter closes by addressing two familiar properties of a matrix: its determinant and rank. Both will be cited throughout this paper.

The rank of an m X n matrix A (denoted by rank(A)) is the number of linearly independent columns in A. Chapter III will show that rank(A) affects the solution of the matrix equation AX = b, where A,X,b are matrices over $\phi(r)$. In finding rank(A), the determinant of a matrix will come into play.

Despite the many techniques which find the determinant of a square matrix, later proofs will refer to the first historical definition of a determinant, computationally inefficient though it is. This definition, later referred to as the "permutational" form of a determinant, derives its name from its indexing (of matrix elements) on a permutation of a set [26:89].

Given the set of integers $S = \{1, 2, ..., N\}$ a permutation on S is a one-to-one and onto mapping σ from S into itself, and is represented by a $2 \times N$ matrix whose top row is S and whose bottom row is a specific permutation of S. For instance, the two permutations of $\{1, 2\}$ are given by

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$$

For $S = \{1, 2, 3, 4\}$ one permutation is given by

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 3 & 1 \end{pmatrix}$$

(where f(1) = 2, f(2) = 4, f(3) = 3 and f(4) = 1) and another by

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix}$$

where f(1) = 4, f(2) = 3, f(3) = 2 and f(4) = 1.

If $A = [a_{k,l}]$ is an N X N matrix, then the determinant of A is given by

$$det(A) = \sum \{sgn(p) | \pi_{j=1,n}(a_{j,p(j)})\}$$
 (2.13)
p(S)

where

S is the set of all permutations on $\{1,2,...N\}$ and

 $sgn(p) = \pm 1$, depending on the number of transpositions which comprise the particular permutation p [7:47-48].

III Solving the Linear Equation AX = b

The key to solving the Lypanov and Riccati equations involves finding the solutions of the linear equation

$$AX = b \tag{3.1}$$

As before, matrices may be non-square but entries must be in $\phi(\tilde{r})$.

Though the matrices of (3.1) may be rectangular, one is first led to investigating properties of a square matrix $A \notin \P^{n\times n}(\Gamma)$ —in particular, whether or not A is invertible. As in the real case, if $\det(A)$ equals zero, then A has no inverse. In general however, the determinant of A is a polynomial over Γ and thus has an inverse except at the roots of $\det(A)$. For example, if

$$A = \begin{bmatrix} x + y & x \\ y & xy \end{bmatrix} \in \mathfrak{t}^{2\times 2}(x,y)$$
 (3.2)

then

det(A) = (x + y)xy - xy = xy(x + y - 1)and A has an inverse except in the root set of det(A)

$$\{(x,y) : x = 0 \text{ or } y = 0 \text{ or } x + y = 1\}$$

If A must be invertible for all values of Γ (as will soon be required), then det(A) can only be a non-zero complex number. In this case A^{-1} is easily found using the well known formula [26:96]

 $A^{-1} = |A|^{-1}$ [matrix-of-cofactors of A]^T

That $A^{-1} \in \P^{n \times n}(\mathbb{R})$ can be seen by recalling that the cofactor of an entry $a_{i,j}$ of A is the determinant of a 3.1

submatrix of A (obtained by deleting the i'th row and j'th column of A). Because the entries of A are in $\mathfrak{C}(\Gamma)$, the cofactors are also in $\mathfrak{C}(\Gamma)$, and thus the transpose of the matrix of cofactors is in $\mathfrak{C}^{n\times n}(\Gamma)$. Since $\det(A)$ is assumed to be a non-zero complex number, so too will A^{l-1} which implies $A^{-1} \in \mathfrak{C}^{n\times n}(\Gamma)$. With this in mind, a specific form of (3.1) can be solved for X: if $A \in \mathfrak{C}^{n\times n}(\Gamma)$ and $\det(A) \in \mathfrak{C} = \{0\}$ then there exists an $A^{-1} \in \mathfrak{C}^{n\times n}(\Gamma)$ and $X = A^{-1}b$ (the general setting for the familiar result involving real matrices). Finally, since the entries of A^{-1} and b are in $\mathfrak{C}(\Gamma)$, so too are the entries in their product $X = A^{-1}b$ as mandated in (3.1).

The picture becomes more complicated when A doesn't have an inverse. This may be because det(A) = 0, $det(A) \notin C(C) - C$, or det(A) isn't defined, i.e. A rectangular. Nevertheless, (3.1) may still have a solution and in fact, may have infinitely many. However, before the general solution of (3.1) can be derived, a more universal setting for 'inverses' of a matrix will be addressed.

The basic concept here involves elementary row and column operations on a matrix $A \in \mathfrak{t}^{m \times n}(P)$. Similar to their counterparts for real matrices, elementary row (column) operations on A are limited to:

interchanging two rows (columns)

n ,

- adding a polynomial multiple of a row (column) to another row (column)
- multiplying a row (column) by a complex scalar not equal to zero.

As with real matrices, an elementary row (column) operation on A (column) is equivalent to left (right) multiplication of A by the corresponding elementary row (column) matrix. For example, interchanging the rows of (3.2) can be accomplished by multiplying A on the left by

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

obtained by interchanging the rows of $I_{2\times2}$. Another example: multiplying the first column of (3.2) by x^2 and adding it to the second column can be accomplished by multiplying A on the right by

$$\begin{bmatrix} 1 & x^2 \\ 0 & 1 \end{bmatrix}$$

obtained by multiplying the first column of $I_{2\times2}$ by x^2 and adding it to the second column. Again, as with real matrices, the determinant of an elementary matrix over $\varphi(r)$ is a nonzero complex number, and thus the elementary matrix always has an inverse. It is for this reason that the third elementary operation is limited to multiplication by a scalar for if polynomial multiples were allowed, the determinant of the resulting elementary matrix may be a polynomial with degree 2 1. The matrix would then fail to have an inverse at the roots of its determinant. A square matrix whose determinant equals 2 1 is called unimodular, and so is invertible.

Both elementary row and column operations are used on A to find the general solution to (3.1). Though two different

cases must be considered -- A does and doesn't have constant rank--both follow the same approach: find unimodular matrices P and Q that reduce A to a form yielding the general solution of (3.1). The two cases will be analyzed separately.

AX = b, rank(A) constant. The aim here is to transform A (¢m×n, using elementary operations, into the m X n matrix

$$A_{1} = \begin{bmatrix} I_{p \times p} & O_{p \times n - p} \\ O_{m - p \cdot p} & O_{m - p \cdot p - p} \end{bmatrix}$$
 (3.3)

where p is the rank of A. The main question thus becomes, "What are the unimodular matrices P and Q such that $PAQ = A_1?$ " Once P and Q are found, the general solution of (3.1) will follow, as this chapter will show. That such P and Q do exist was recently proven by E. Sontag [31]:

<u>THEOREM 3.0 (Sontag)</u>: The following statements are equivalent for a matrix A = A(r) over $R = \phi(r)$:

- There exists a matrix B over R such that ABA = A and BAB = B. B is called the weak generalized inverse (or the {1,2}-generalized inverse) of A.
- ь. There exists square unimodular matrices P and Q over R such that A: = PAQ.
- As a function of the the complex variables $\Gamma = (\Gamma_1, \dots, \Gamma_n)$, rank(A) is constant.

Proof: See [31].

A method for determining P and Q has been developed by Dr. John Jones Jr., AFIT, and works by keeping a 'cumulative' record of all the elementary row and column operations performed in reducing A to A:. The "Jones ST

Method" begins by forming the matrix

$$A_{z} = \begin{bmatrix} \frac{A_{m \times n}}{I_{n \times n}} & I_{m \times m} \\ I_{n \times n} & O_{n \times m} \end{bmatrix}$$
 (3.4)

Subsequent elementary operations are done on A_2 until A becomes A_1 . Once this point is reached, the matrix occupying the $I_{m\times m}$ block is P, and the matrix occupying the $I_{m\times m}$ block is Q. That these two blocks do indeed contain P and Q is easily seen: P and Q will be the respective products of the elementary row and column transformations done in reducing A to A_1 ; since $I_{m\times m}$ is row adjacent to A, P will be the left multiplier of A, and since $I_{n\times m}$ is column adjacent to A, Q will be A's right multiplier. Also, both P and Q are invertible, since their determinants are nonzero complex. To see this, consider $P = \pi R_1$ where the R_1 's are the elementary row transformations done in reducing A to A_1 . It follows that

$$det(P) = det(\pi R_1) = \pi det(R_1)$$

Since each $det(R_1)$ is nonzero complex (the nature of an elementary matrix), so will their product be, and thus det(P). A similar approach holds for Q.

Example: Reduce to the form of (3.3) the matrix

$$A = \begin{bmatrix} 1 & x & y \\ 0 & 0 & 1 \end{bmatrix} \quad \xi \quad \varphi^{2\times 3}(x,y)$$

Begin by augmenting A by the identity matrices $I_{2\times2}$ and $I_{3\times3}$ to form per (3.4)

The next step is to get an identity matrix in the upper left hand corner of A, per (3.3). First, interchange the second and third columns to form

0	y 1	х О	1 0	0
1	0	0		
0	0	1	03,	(2
0	1	0		

Next, add -y times the second row and add it to the first to form

Next, add -x times the first column to the third column obtaining

The process is now complete since A has been reduced to the form of (3.3). The P and Q elementary row and column matrices are read off to be

$$P = \begin{bmatrix} 1 & -y \\ 0 & 1 \end{bmatrix} \qquad \text{and} \qquad Q = \begin{bmatrix} 1 & 0 & -x \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

Thus

$$PAQ = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Since det(P) = 1 and det(Q) = -1, P and Q are unimodular and hence invertible.

The above example illustrates a theorem helping to identify the general solution of (3.1) for A of constant rank:

THEOREM 3.1: Let A f $f^{m \times n}(n)$ and rank(A) = p a constant. If P and Q are unimodular and

$$PAQ = \begin{bmatrix} I_{PXP} & O_{PXP-P} \\ O_{m-PXP} & O_{m-PXP-P} \end{bmatrix}$$

where

$$P = \prod_{M} \{ \varphi_{m \times m}(\Gamma) , Q = [S \ N] \{ \varphi_{n \times n}(\Gamma) \},$$

$$T \{ \varphi_{n \times m}(\Gamma) , M \{ \varphi_{m - P}, m(\Gamma) \}, S \{ \varphi_{n \times P}(\Gamma) \},$$

$$N \{ \varphi_{n \times n - P}(\Gamma) \}$$

then

$$\begin{bmatrix} A & I_{m\times m} \\ I_{D\times D} & O_{D\times m} \end{bmatrix} \text{ is similar to } \begin{bmatrix} I_{p\times p} & O_{p\times D-p} & T_{p\times m} \\ O_{m-p\times p} & O_{m-p\times D-p} & M_{m-p\times m} \\ S_{D\times p} & N_{D\times D-p} & O_{D\times m} \end{bmatrix}$$

$$PROOF: \begin{bmatrix} P_{m\times m} & O_{m\times D} \\ O_{D\times m} & I_{D\times D} \end{bmatrix} \begin{bmatrix} A_{m\times D} & I_{m\times m} \\ I_{D\times D} & O_{D\times m} \end{bmatrix} \begin{bmatrix} Q_{D\times D} & O_{D\times m} \\ O_{m\times D} & I_{m\times m} \end{bmatrix}$$

$$= \begin{bmatrix} (PAQ)_{m\times D} & P_{m\times m} \\ I_{D\times D} & O_{D\times m} \end{bmatrix} \begin{bmatrix} Q_{D\times D} & O_{D\times m} \\ O_{m\times D} & I_{m\times m} \end{bmatrix}$$

$$= \begin{bmatrix} (PAQ)_{m\times D} & P_{m\times m} \\ Q_{D\times D} & O_{D\times m} \end{bmatrix} \text{ (from given and)}$$

$$= \begin{bmatrix} (PAQ)_{m\times D} & P_{m\times m} \\ Q_{D\times D} & O_{D\times m} \end{bmatrix} \text{ (substitute for P & Q)}$$

$$= \begin{bmatrix} I_{p\times p} & O_{p\times n-p} & T_{p\times n} \\ O_{m-p\times p} & O_{m-p\times n-p} & M_{m-p\times m} \\ \end{bmatrix}$$

Since P and Q are unimodular,

$$\begin{bmatrix} P_{m\times m} & O_{m\times n} \\ O_{n\times m} & I_{n\times n} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} Q_{n\times n} & O_{n\times m} \\ O_{m\times n} & I_{m\times m} \end{bmatrix}$$

are also unimodular and thus invertible. This is seen by noting that if A and B are square matrices, then

$$\det \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} = \det(A) * \det(B)$$

a consequence of the permutational definition of a determinant (2.13).

Example of Theorem 3.1: Let

0.

$$A = \begin{bmatrix} x & y & x^2 + y + 1 \\ x^2y & xy^2 & x^3y + xy^2 + xy \end{bmatrix}$$

Matrices P and Q are sought that reduce A to the form (3.3). First, augment A by the identity matrices $I_{2\times 2}$ and $I_{3\times 3}$ to get

Next, add -xy times row 1 and add to row 2 to obtain:

Next, add -1 times column 2 to column 3 to get:

× 0	у 0	x² + 1 0	1 -xy	0
1	0	0		
0	1	-1	03	×2
0	0	1	ļ	

Now add -x times column 1 to column 3 to obtain:

× o	ý	0	1 -xy	0
1	0	-x		
0	1	-1	03	x2
0	0	1	1	

Switching columns 1 and 3:

0	у О	х 0	1 -×y	0
	0	1		
-1	1	0	03	x2
1	0	0		

Adding -y times column 1 to column 2, followed by -x times column 1 to column 3 yields:

1 0	0 0	[1 [-xy	0] >> T 1] >> M
-x -1	xy 1 + x ² 1 + y x	03×2	-
i	-y , -x	V3X2	
^	\ /		
S	N		

Since A has been reduced to the form (3.3), P and Q have been found, and thus S,T,M,N of Theorem 3.1. It's easy to verify the unimodularity of P and Q.

The groundwork is now in place to identify the general form of the solution to (3.1). The following theorem is due to Jones [11]:

THEOREM 3.2: Let A ($\phi^{m\times n}(\Gamma)$ of constant rank p and b ($\phi^{m\times n}(\Gamma)$). Then AX = b has a solution X ($\phi^{n\times n}(\Gamma)$) if and only if Mb = $O_{m-p\times n}$, M given in Theorem 3.1. The general form of X is given by

$$X = STb + NZ$$

S,T,N as given in Theorem 3.1, and Z ($^{n-p}$ $^{\vee}$ (P) arbitrary.

PROOF: Let P ($\phi^{m\times m}(\Gamma)$ and Q ($\phi^{n\times n}(\Gamma)$ be unimodular matrices such that

$$PAQ = \begin{bmatrix} I_{pxp} & O_{pxn-p} \\ O_{m-p,p} & O_{m-p,n-p} \end{bmatrix}$$

as given in Theorem 3.0. Then AX = b has a solution X iff

PAX = Pb has a solution X (by virture of the unimodularity of P and hence the existence of P^{-1})

 $PA(QQ^{-1})X = Pb = (PAQ)(Q^{-1}X)$ has a solution X (since Q is unimodular, there exists Q^{-1})

$$(PAQ)Y = Pb$$
 has a solution $Y = Q^{-1}X$ (3.5)

Let

$$Y = \begin{bmatrix} W_{PXV} \\ Z_{D-PXV} \end{bmatrix}$$
, $P = \begin{bmatrix} T_{PXM} \\ M_{M-PXM} \end{bmatrix}$, $Q = [S_{DXP} \ N_{DXD-P}]$

Then substituting for PAQ, P and Y in (3.5):

$$\begin{bmatrix} I_{PXP} & O_{PXP-P} \\ O_{m-P,P} & O_{m-P,P-P} \end{bmatrix} \begin{bmatrix} W \\ Z \end{bmatrix} = \begin{bmatrix} T \\ M \end{bmatrix} b$$
$$\begin{bmatrix} W \\ O_{m-PXV} \end{bmatrix} = \begin{bmatrix} Tb \\ Mb \end{bmatrix}$$
(3.6)

Since $Y = Q^{-1}X$ from (3.5),

$$X = QY = [S \ N] \begin{bmatrix} W \\ Z \end{bmatrix} = SW + NZ = STb + NZ$$

Note that (3.6) allows Z to be arbitrary since Z is zeroed out by the second row of PAQ. (3.6) also says that if a

solution exists to AX = b , then Mb must equal $O_{m-p \times v}$. This mandate is referred to as the 'consistency condition'.

EXAMPLE (THEOREM 3.2):

Given

$$A = \begin{bmatrix} x & y & x^{2} + y + 1 \\ x^{2}y & xy^{2} & x^{3}y + xy^{2} + xy \end{bmatrix}$$

$$b = \begin{bmatrix} x + xy + y + x^{2}y + y^{2} \\ x^{2}y + xy^{2} + x^{2}y^{2} + x^{3}y^{2} + xy^{3} \end{bmatrix}$$

solve AX = b.

The example for Theorem 3.1 (which reduced A) found that

 $T = [1 \ 0]$, $M = [-xy \ 1]$, $S = [-x \ -1 \ 1]^c$ and

$$N = \begin{bmatrix} xy & 1 + x^2 \\ 1 + y & x \\ -y & -x \end{bmatrix}$$

It is a simple matter to show Mb = 0 thereby confirming the existence of a solution for AX = b. From Theorem 3.2 it follows that the general solution is

$$X = STb + NZ =$$

$$\begin{bmatrix} -x^{2} - x^{2}y - xy - x^{3}y - xy^{2} \\ -x - xy - y - x^{2}y - y^{2} \\ x + xy + y + x^{2}y + y^{2} \end{bmatrix} + a \begin{bmatrix} xy \\ 1 + y \\ -y \end{bmatrix} + \beta \begin{bmatrix} 1 + x^{2} \\ x \\ -x \end{bmatrix}$$
where a, \beta \infty \infty (x,y)

A benefit of Theorem 3.2 is that it easily identifies the basis for the kernal of the linear transformation represented by the m X n matrix A, a task generally quite tedious. Recall that the kernal of a linear transformation 3.11

represented by the matrix A is the set

$$K_A = \{ X : AX = O_{m \times 1} \}$$

To find the basis of K_{\triangle} , one must solve the equation

$$AX = O_{m \times 1}$$

which is a special case of AX = b where $b = 0_{m \times 1}$ From Theorem 3.2, it follows that

$$X = STb + NZ = STO_{mx1} + NZ = NZ$$

Thus the span of the columns of N is Ka. However, since $Q = [S \ N]$ and Q is unimodular, the columns of Q are linearly independent, and thus the columns of N. Since the columns of N is a linearly independent spanning set of K_A , it follows that the columns of N are the basis for the kernal of the linear transformation represented by the matrix A.

AX = b, rank(A) not constant. Consider the following matrix $A \in \mathcal{C}^{3\times 3}(x,y,z)$ which doesn't have constant rank:

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$$
 (3.7)

Rank(A) $\{$ {0,1,2,3} , depending on the values assumed by (x,y,z): if x = y = z = 0, then rank(A) = 0; if $x,y,z \neq 0$, then rank(A) = 3; if two are 0 while the third isn't, rank(A) = 1; if two aren't zero while the third is, then rank(A) = 2. This section addresses the solution of the equation AX = b given rank(A) not constant.

A result similar to Theorem 3.2 holds for such a matrix A. The approach is practically identical to the

constant rank case: find unimodular matrices P and Q which reduce A to the form

$$PAQ = \begin{bmatrix} I_{p \times p} & O_{p \times n - p} \\ O_{m - p \times p} & Q_{m - p \times n - p} \end{bmatrix}, p \ge 1$$
 (3.8)

where p is as large as possible and $\Omega \in C^{m-p\times n-p}(P)$. As before, P and Q are found using the Jones ST Method. If such P and Q are found (they may not exist), the following theorem applies:

THEOREM 3.3: Let A (cm×n(r) If P and Q are unimodular and

$$PAQ = \begin{bmatrix} I_{PXP} & O_{PXP-P} \\ O_{m-PXP} & A_{m-PXP-P} \end{bmatrix}$$

where

 $P = \begin{bmatrix} T \\ M \end{bmatrix} \quad \text{$($ $\varphi_{xm}(\Gamma)$ }, \quad Q = [S \ N] \quad \text{$($ $\varphi_{xm}(\Gamma)$ }, \\ T \quad \text{$($ $\varphi_{xm}(\Gamma)$ }, \quad M \quad \text{$($ $\varphi_{m-p+m}(\Gamma)$ }, \quad S \quad \text{$($ $\varphi_{nxp}(\Gamma)$ }, \\ N \quad \text{$($ $\varphi_{nxn-p}(\Gamma)$ }, \quad \Omega \quad \text{$($ $\varphi_{m-pxn-p}(\Gamma)$ }) \\ \text{then}$

$$\begin{bmatrix} A & I_{m\times m} \\ I_{n\times n} & O_{n\times m} \end{bmatrix} \quad \text{is similar to} \quad \begin{bmatrix} I_{p\times p} & O_{p\times n-p} & T_{p\times m} \\ O_{m-p\times p} & I_{m-p\times n-p} & M_{n-p\times m} \\ S_{n\times p} & N_{n\times n-p} & O_{n\times m} \end{bmatrix}$$

PROOF: identical to that of Theorem 3.1. ■

Once P and Q have been found, the following theorem is applied:

THEOREM 3.4: Let $A \in \mathfrak{t}^{m \times n}(\Gamma)$ and $b \in \mathfrak{t}^{m \times n}(\Gamma)$. Then AX = b has a solution $X \in \mathfrak{t}^{n \times n}(\Gamma)$ if and only if there exists a $Z \in \mathfrak{t}^{n-p} \cdot r(\Gamma)$ such that $Mb = \Lambda Z = M$, Λ given in Theorem 3.3. In this case, the general form of X is given

$$X = STb + NZ$$

for all Z such that Mb = AZ and S,T,N given in Theorem 3.3.

PROOF: Let P $\{ \phi^{m\times m}(r) \}$ and $\mathbb{Q} \{ \phi^{n\times n}(r) \}$ be

unimodular matrices such that

$$PAQ = \begin{bmatrix} I_{PXP} & O_{PXP-P} \\ O_{m-P,P} & A_{m-P,P-P} \end{bmatrix}$$

AX = b has a solution X

iff

PAX = Pb has a solution X (by virture of the unimodularity of P and hence the existence of A^{-1})

 $PA(QQ^{-1})X = Pb = (PAQ)(Q^{-1}X)$ has a solution X (since Q is unimodular, there exists Q^{-1})

$$(PAQ)Y = Pb \text{ has a solution } Y = Q^{-1}X$$
 (3.9)

Let

$$Y = \begin{bmatrix} W_{PXV} \\ Z_{D-PXV} \end{bmatrix}, P = \begin{bmatrix} T_{PXM} \\ M_{M-PXM} \end{bmatrix}$$

and

$$Q = [S_{n \times p} \ N_{n \times n - p}]$$

Then substituting for PAQ, P and Y in (3.9):

$$\begin{bmatrix} I_{PXP} & O_{PXP-P} \\ O_{M-P,P} & \Omega_{M-P,P-P} \end{bmatrix} \begin{bmatrix} W \\ Z \end{bmatrix} = \begin{bmatrix} T \\ M \end{bmatrix} b$$

$$\begin{bmatrix} W \\ \Omega Z \end{bmatrix} = \begin{bmatrix} Tb \\ Mb \end{bmatrix}$$
(3.10)

Since $Y = Q^{-1}X$.

$$X = QY = [S \ N] \begin{bmatrix} W \\ Z \end{bmatrix} = SW + NZ = STb + NZ$$

subject to

$$\Omega Z = Mb \tag{3.11}$$

per (3.10). This latter mandate is the consistency

condition. That Z ($\phi^{n-p\times \vee}(P)$ proceeds from (3.9): since 3.14

Q has polynomial entries and is unimodular, so too is Q^{-1} . Since X is also required to be in $\Phi^{n\times \vee}(P)$ the product of Q^{-1} and X must be in $\Phi^{n\times \vee}(P)$. Since Z is a submatrix of $Y = Q^{-1}X$, Z $\Phi^{n-p\times \vee}(P)$.

In general, there is no assurance that the consistency condition (3.11) can be met for a given equation AX = b: though A may be reduced to the form (3.8), there may not exist a Z OVER (Γ) satisfying the consistency condition AZ = Mb. It may be necessary to leave (Γ) in order to find a Z satisfying the consistency condition, an act contrary to this paper's aim.

What then can be said about the solvability over $\mathfrak{t}(n)$ of AX = b when rank(A) isn't constant? It's possible that A isn't reducible to the form (3.8). For instance the matrix in (3.7): reducing A to the form (3.8) implies that A has a rank no less than p (the first p columns of (3.8) are linearly independent despite the form of \mathfrak{Q}). However, the rank of A may be zero if x = y = z = 0, and so A isn't reducible to the form (3.8).

This example, however, is symptomatic of a larger problem. Assume that a given matrix A has been reduced as much as possible (no known method exists that insures maximum reducibility of a matrix over $\Phi(P)$, P arbitrary), i.e. the maximum value of p has been found. One is then left with finding a Z satisfying the consistency condition. A in turn cannot be reduced, since this would violate the

assumption that A has been reduced as far as is possible. If Ω is square, $\det(\Omega)$ cannot be a nonzero complex number (if it were, $\operatorname{rank}(\Omega) = m - p$ and Ω could be reduced to $\operatorname{I}_{m \sim p \times m - p}$, thus giving A constant rank, a contradiction). If $\det(\Omega) \neq 0$ repeated use of Cramer's rule may identify some solutions Z, but these would be local since the roots of $\det(\Omega)$ must be omitted. In general, however, Ω will be rectangular.

Thus finding the general solution of (3.1) given rank(A) not constant ultimately requires solving (3.11) where Ω cannot be transformed into the form (3.8). The search for Z may be aided by matrices related to Ω , the topic of the next section.

SMITH FORM OF A MATRIX. A matrix A (r) is said to be in Smith Form (SF) if and only if

$$A = \begin{bmatrix} f_1 & 0 & & & \\ & & & & \\ 0 & f_3 & & & \\ \hline 0_{m-1}x_1 & & 0_{m-1}x_{m-1} \end{bmatrix}$$
 (3.12)

where all off diagonal entries equal zero and f_{κ} is a factor of $f_{\kappa+1}$. For example

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & x^{2}y & 0 \\ 0 & 0 & x^{2}y^{2} \end{bmatrix} \quad (x^{3}x^{3}(x,y))$$

is in SF (3.12) because all off diagonal entries are zero, x is a factor of x^2y , and x^2y is a factor of x^2y^3 . Another example is

$$A = \begin{bmatrix} x & 0 \\ 0 & x(x+1) \\ 0 & 0 \end{bmatrix} \quad (x^{3\times2}(x,y))$$

which is in SF (3.12) because all off diagonal entries are zero, and x is a factor of x(x+1). The solving of matrix equation (3.1) may be aided if A can be transformed (using unimodular matrices P' and Q') into a matrix having SF. In fact, Smith Forms have already appeared: (3.3) is a special case where $f_k=1$, $k \le p$ The main aim here is to use SFs to solve the equation (3.11) $\Omega Z=Mb$ of the last section.

To illustrate, assume that in solving $\Omega Z = Mb$ (derived from some hypothetical original equation AX = b), unimodular matrices P' and Q' have been found that give P' ΩQ ' a SF. Thus

$$\Omega Z = Mb$$

 $P'\Omega Z = P'Mb$
 $P'\Omega(Q'Q'^{-1})Z = P'Mb$
 $(P'\Omega Q')(Q'^{-1}Z) = P'Mb$

Letting

$$Y = Q'^{-1}Z$$

$$H = P'Mb$$

$$F = P'\Omega Q'$$
(3.13)

the last equality becomes

$$FY = H$$

Since F is in SF, there exists at most one nonzero entry in each row and column of F. Because F and H are known, some entries in Y can be found by dividing the respective F and H factors (if however a given entry of Y isn't compatible with two or more equations, or if the entry doesn't turn out to 3.17

be a polynomial, then $\Omega Z = Mb$ has no solution Z.) Depending on the structure of F, there may be entries of Y which can assume arbitrary values. Y is then substituted into (3.13) to yield Z = Q'Y, thus solving (3.11) and in turn (3.1).

Given A ($t^{m\times n}(\Gamma)$, if there exists unimodular matrices P' ($t^{m\times m}(\Gamma)$ and Q' ($t^{n\times n}(\Gamma)$) such that A' = P'AQ' is in SF, then A is said to be equivalent to its Smith Form. It has been proven [1:188] that for every A ($t^{m\times n}(x)$), A is equivalent to a unique Smith Form (an algorithm exists [1:192] which finds this). However, such is not always the case for multi-dimensional Γ . In fact, Lee and Zak [20] prove that a matrix A ($t^{m\times n}(x,y)$ is equivalent to its SF iff a certain system of linear polynomial equations has a solution. The conditions under which a matrix in three or more variables is equivalent to its Smith Form remains an open question.

Although the existence of a SF for a given matrix A must be known before a search for it can begin, a method is needed which identifies the unimodular P' and Q' which yield a particular SF of A. As before, the Jones ST Method can be used.

EXAMPLE: Find the Smith Form of the matrix (Frost and Storey [8])

$$A = \begin{bmatrix} 5 + z & 0 & 1 \\ 0 & 5 + z & 0 \\ 0 & 0 & 5 \end{bmatrix} \quad (\ ^{3\times 3}(5,z))$$

Begin by augmenting A with Isxs:

Next, add -(z + s) times the third column to the first column to get:

Add -s times row 1 to row 3 obtaining:

Finally, interchange the first and third columns:

Since the upper left hand matrix is in Smith Form (i.e., 1 is a factor of s + z, in turn a factor of -s(s + z) and off diagonal entries are zero)

$$P' = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -s & 0 & 1 \end{bmatrix} \qquad Q' = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 - (s+z) \end{bmatrix}$$

are the two unimodular (easily verifiable) matrices which reduce A to its Smith Form.

IV Solutions to Lypanov and Riccati Equations

This chapter will consider the solutions to the Lypanov matrix equation

$$AX - XB = C (4.1)$$

and the second order Riccati matrix equation

$$XDX + AX + XB + C = 0 (4.2)$$

where the entries of A,B,C,D are in $\phi(n)$. It will be shown that solving these two equations for X hinges upon solving equations of the form AX = b the subject of Chapter III.

Lypanov Equation. (4.1) may be solved using tensor products or similarity transformations. The former approach will be addressed first because it uses the results of Chapter III at the onset. The approach using similarity transformations will then be presented as a lead-in to solving the Riccati equation (4.2), whose solution relies exclusively on these tranformations. Whichever method is taken though, the linear equation (3.1) will demand resolution.

Lypanov Equation: Tensor Solution. Multiplication between two matrices is easily extended to that of their tensor product. Given $A = \begin{bmatrix} a_1 \end{bmatrix}$ the tensor product of the matrices A and B, represented by AAB, is given by $\begin{bmatrix} a_1 \end{bmatrix}$. Unlike matrix multiplication, the number of columns in A doesn't have to equal the number of rows in B.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \qquad \text{and} \qquad B = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$$

then

It follows that if A is m X n and B is r X s, then $A \triangle B$ is mr X ns.

The next step in solving (4.1) is to identify the solution of the equation

$$AXB = C (4.3)$$

where $A_{i,r}$ denotes the i'th row of A and $A_{i,c}$ its i'th column (the i-jth entry of A is given by $a_{i,j}$ with no comma between subscripts).

THEOREM 4.1: Solving the matrix equation

is equivalent to solving the equation

where

$$G = AAB^T$$
, $u = [X_1, \dots, X_m, r]^T$, $c = [C_1, \dots, C_m, r]^T$
 $PROOF$: Let $A = [a_{x,x}]$, $B = [B_{x,x}]$,
 $C = [c_{x,x}]$, $X = [x_{x,x}]$

Since AXB = C,

CIJ = AI, - XBJ, - = AI, - [X1, -BJ, - ... Xn, -BJ, -] T = Ek[aIk EL (XkLBLJ)] = Ek, L (aIkBLJX kL) = EL (aIIBLJX 1L) + *** + EL (aInBLJX nL) = aII EL (BLJX 1L) + *** + aIn EL (BLJX nL) = [aIIBJJ...BpJ] ... aIn[BJJ...BpJ]] * [X1, -... Xn, -] T = [aIIBJJ... aInBJJ...] * [X1, -... Xn, -] T (4.4)

Since $C = [C_1, \dots, C_m, -]^T$ is given and from (4.4), the theorem follows.

The following theorem is due to P. Lancaster [18]:

Theorem 4.2: Given the matrix equation

$$A_1XB_1 + A_2XB_2 = C \tag{4.5}$$

where A_{x} are m X m , B_{x} are n X n , X,C are m X n (with X_{x} and C_{x} the rows of X and C respectively), (4.5) is equivalent to solving the equation $G_{x} = c$ where

$$G = A_1 \triangle B_1^{\mathsf{T}} + A_2 \triangle B_2^{\mathsf{T}}$$

$$\times = [X_1 \dots X_m]^{\mathsf{T}}$$

$$C = [C_1 \dots C_m]^{\mathsf{T}}$$

Proof: Similar to that of Theorem 4.1.

The restriction on square A_x and B_x in Theorem 4.2 is easily modified to include rectangular matrices. With this modification, Theorem 4.2 allows for the solution of the Lypanov equation (4.1) which is a special case of (4.5) with $B_1 = I$ and $A_2 = -I$. Once a given Lypanov equation has been reduced to an equation of the form AX = b using tensor products, the results of Chapter III can be applied to this latter equation.

Example: Solve the Lypanov equation

$$AX + XB = C = AXI + IXB$$
4.3

where

$$A = \begin{bmatrix} s+z & s \\ z & s-z \end{bmatrix}, \quad B = \begin{bmatrix} -z & -s \\ -z-s & s-z \end{bmatrix}$$

$$C = \begin{bmatrix} sz & s^2 \\ 2s^2-2z^2-sz & 4sz-2z^2-3s^2 \end{bmatrix}$$

$$X = \begin{bmatrix} y & z \\ y & w \end{bmatrix}$$
(4.6)

The first step is to reduce the given Lypanov equation to the form Gx = c so that Theorem 4.2 may be used. Thus $G = A \blacktriangle I^\intercal + I \blacktriangle B^\intercal =$

$$\begin{bmatrix} s+z & s \\ z & s-z \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} -z & -z-s \\ -s & s-z \end{bmatrix} = \begin{bmatrix} s+z & 0 & s & 0 \\ 0 & s+z & 0 & s \\ z & 0 & s-z & 0 \\ 0 & z & 0 & s-z \end{bmatrix} + \begin{bmatrix} -z & -z-s & 0 & 0 \\ -s & s-z & 0 & 0 \\ 0 & 0 & -z & -z-s \\ 0 & 0 & -s & s-z \end{bmatrix} = \begin{bmatrix} s & -z-s & s & 0 \\ -s & 2s & 0 & s \\ z & 0 & s-2z & -z-s \\ 0 & z & -s & -2s-2z \end{bmatrix}$$

Let $x = [y z v w]^T$ and

 $c = [sz s^2 2s^2-2z^2-sz 4sz-2z^2-3s^2]^T$. Thus the solution

$$X = \begin{bmatrix} y & z \\ & & \end{bmatrix}$$

to the given Lypanov equation is the solution to the linear equation

$$\begin{bmatrix} s & -z-s & s & 0 \\ -s & 2s & 0 & s \\ z & 0 & s-2z & -z-s \\ 0 & z & -s & -2s-2z \end{bmatrix} \begin{bmatrix} y \\ z \\ v \\ w \end{bmatrix} = \begin{bmatrix} sz \\ s^2 \\ 2s^2-2z^2-sz \\ 4sz-2z^2-3s^2 \end{bmatrix}$$

Depending on the nature of the G matrix obtained using tensor products, the given Lypanov equation may or may not be easily solvable. For instance, if G cannot be reduced to the form (3.3), one then tries for form (3.8). Here however, finding the Z's satisfying the consistency condition (3.11) $\Omega Z = Mb$ may be difficult. In any event, the use of tensor products creates a large matrix G, thereby compounding calculations.

The second approach to solving the Lypanov equation

(4.1) uses similarity transformations. Although this method may not explicitly identify the solution to (4.1), it will identify a set of matrices within which the solutions of (4.1) will be found.

$$E = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix}$$
 (4.7)

always has det(E) = 1 despite the nature of X (this follows from the permutational definition of a determinant (2.13). Since E is unimodular, it has the unimodular inverse

$$E' = \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix} \tag{4.8}$$

which is verified below:

$$\mathsf{EE.} = \begin{bmatrix} 0 & \mathbf{I} \\ \mathbf{I} & \mathbf{X} \end{bmatrix} \begin{bmatrix} 0 & \mathbf{I} \\ \mathbf{I} & -\mathbf{X} \end{bmatrix} = \begin{bmatrix} 0\mathbf{I} + \mathbf{I}\mathbf{0} & -0\mathbf{X} + \mathbf{I}\mathbf{5} \\ \mathbf{I} + \mathbf{X}\mathbf{0} & -\mathbf{I}\mathbf{X} + \mathbf{X}\mathbf{I} \end{bmatrix} = \begin{bmatrix} 0 & \mathbf{I} \\ \mathbf{I} & \mathbf{0} \end{bmatrix}$$

 $(E'E = I_{2n\times 2n} \text{ likewise follows}).$

The next observation uses the matrix

$$R = \begin{bmatrix} A & C \\ O & B \end{bmatrix} \tag{4.9}$$

where A,B,C are all nXn matrices:

$$ERE^{-1} = \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} A & C \\ O & B \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix} = \begin{bmatrix} A & C+XB \\ O & B \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix}$$
$$= \begin{bmatrix} A & -AX + C + XB \\ O & B \end{bmatrix}$$
(4.10)

The above two observations can now be brought together to solve (4.1): the 1-2'th entry of ERE^{-1} above is itself a Lypanov equation. If it equalled zero, then X would be a solution to the equation

$$-AX + XB + C = O$$
 or $AX - XB = C$

which is of the form (4.1). The following theorem due to Jones [11] has now been proven:

<u>Theorem 4.3</u>: If X is a solution to the Lypanov equation

$$AX - XB = C$$

where A,B,C,X (¢°×°(°) then

$$R = \begin{bmatrix} A & C \\ O & B \end{bmatrix}$$
 is similar to
$$R' = \begin{bmatrix} A & O \\ O & B \end{bmatrix}$$
 (4.11)

Proof: previous discussion.

The following more powerful theorem was proven by Roth [30] for matrices whose entries are complex numbers:

Theorem 4.4: If A,B,C ($\phi^{n\times n}$, then AX - XB = C has a solution X ($\phi^{n\times n}$ if and only if R and R' in (4.11) are similar.

Proof: See [30]. ■

Although Theorem 4.3 is the core result leading to a solution of (4.1), the final assault upon (4.1) will use polynomials of a matrix. For this reason, the following theorems are presented.

Theorem 4.5: Let A,B be n X n matrices such that B^{-1} exists. Then for n \in Z+

$$(BAB^{-1})^n = BA^nB^{-1}$$
 (4.12)

Proof: Inductive reasoning will be used. (4.12) is trivially true for n = 1. For n = 2,

 $(BAB^{-1})^2 = (BAB^{-1})(BAB^{-1}) = BA(B^{-1}B)AB^{-1}$ = BA(I)AB⁻¹ = BA²B⁻¹ and (4.12) is again true. Assume (4.12) is true for n = k Thus

$$(BAB^{-1})^k = BA^kB^{-1}$$

and so

$$(BAB^{-1})^{k+1} = (BAB^{-1})^{k}(BAB^{-1}) = BA^{k}B^{-1}(BAB^{-1})$$

= $BA^{k}(B^{-1}B)AB^{-1} = BA^{k+1}B^{-1}$

and so (4.12) holds true for k + 1 and therefore for all

positive integers.

Theorem 4.6: Let f(x) be a polynomial in the variable x. Let A,B be n X n matrices such that B^{-1} exists. Then

$$f(BAB^{-1}) = B f(A) B^{-1}$$

Proof: Let $f(x) = a_0 + \sum_{k=1,n} a_k x^k$ Then

$$f(BAB^{-1}) = a_0I + \Sigma a_K (BAB^{-1})^K = (from Theorem 4.5)$$

= $a_0BB^{-1} + \Sigma a_K BA^K B^{-1}$
= $Ba_0B^{-1} + B (\Sigma a_K A^K) B^{-1}$
= $B (a_0I + \Sigma a_K A^K) B^{-1}$
= $B f(A) B^{-1} \blacksquare$

Theorem 4.7: Let

$$R = \begin{bmatrix} A & O \\ B & C \end{bmatrix}$$

A,B,C nXn matrices. Then for K & Z+

$$R^{K} = \begin{bmatrix} A^{K} & 0 \\ * & C^{K} \end{bmatrix}$$
 (4.13)

* an expression in A,B,C.

Proof: Inductive reasoning will be used. (4.13) is trivially true for i = 1, where * = B = OA + 1B + OC

For i = 2,

$$R^{2} = \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} = \begin{bmatrix} A^{2} & 0 \\ BA+CB & C^{2} \end{bmatrix}$$

and (4.13) holds true again. Let (4.13) be true for n=k. Then

$$R^{k+1} = R^k R = \begin{bmatrix} A^k & 0 \\ * & C^k \end{bmatrix} \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} = \begin{bmatrix} A^{k+1} & 0 \\ * & C^{k+1} \end{bmatrix}$$

and thus (4.13) is true for all positive integers.

Theorem 4.8: Let

$$R = \begin{bmatrix} A & O \\ B & C \end{bmatrix}$$

A,B,C nXn matrices. If f(x) is a polynomial in x, then

$$f(R) = \begin{bmatrix} f(A) & 0 \\ + & f(C) \end{bmatrix}$$
 (4.14)

* an expression in A,B,C

Proof: Inductive reasoning will be used. Let $f(x) = a_0 + \sum_{k=1.na_k x^k} For \ n = 1, \ f(R) = a_0 + a_1 R = a_0 \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} + a_1 \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} = \begin{bmatrix} a_0 I + a_1 A & 0 \\ a_1 B & a_0 I + a_1 C \end{bmatrix} = \begin{bmatrix} f(A) & 0 \\ * & f(C) \end{bmatrix}$

where

$$* = a_1B = a_1B + OA + OC$$

and so (4.14) holds.

For n = 2,
$$f(R) = a_0I + a_1R + a_2R^2$$

= $a_0 \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} + a_1 \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} + a_2 \begin{bmatrix} A^2 & 0 \\ * & C^2 \end{bmatrix}$ (from Theorem 4.7)

= $\begin{bmatrix} a_0I + a_1A + a_2A^2 & 0 \\ * & a_0I + a_1C + a_2C^2 \end{bmatrix}$

= $\begin{bmatrix} f(A) & 0 \\ * & f(C) \end{bmatrix}$

and (4.14) remains true. Let (4.14) be true for n = v

If
$$f_1(x) = [a_0 + E_{k=1}, \sqrt{a_k}x^k]$$
 and $f(x) = f_1(x) + a_{v+1}x^{v+1}$ then $f(R) = [f_1(A) \ 0] + a_{v+1}R^{v+1}$

$$= \begin{bmatrix} f_1(A) & 0 \\ * & f_1(C) \end{bmatrix} + a_{v+1}R^{v+1}$$

$$= \begin{bmatrix} f_1(A) & 0 \\ * & f_1(C) \end{bmatrix} + a_{v+1}\begin{bmatrix} A^{v+1} & 0 \\ * & C^{v+1} \end{bmatrix}$$

$$= \begin{bmatrix} f_1(A) & 0 \\ * & f_1(C) \end{bmatrix} + a_{v+1}[A^{v+1}]$$

$$= \begin{bmatrix} f_{1}(A) + a_{\vee+1}A^{\vee+1} & 0 \\ * & f_{1}(C) + a_{\vee+1}C^{\vee+1} \end{bmatrix}$$

$$= \begin{bmatrix} f(A) & 0 \\ * & f(C) \end{bmatrix}$$

and so (4.14) is true for all positive integers.

One more theorem remains to be cited before the assault on (4.1) can begin. The Hamilton-Cayley Theorem will permit the identification of a pair of linear equations whose common solutions will contain those of (4.1).

Theorem 4.9 (Hamilton-Cayley): Given A an nXn matrix and $\mathfrak{F}(y) = \det(A - yI_{n\times n})$ the characteristic equation of A, then

$$\xi(A) = 0_{m \times m}$$

Proof: see Nering [26:100]

The tools are now in place with which to solve (4.1). The following theorem is due to Jones [11]:

Theorem 4.10: Given the Lypanov equation

$$AX - XB = C \quad A_1B_1C \in \mathfrak{C}^{n \times n}(\Gamma) \tag{4.15}$$

let R and R' be defined as in (4.11) and f_{A} the characteristic equation of A. If

$$f_{A}(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
 (4.16)

then

$$U + XV = 0$$
 (4.17)
M + XN = 0

Thus a solution to (4.15) will be found among the common solutions of the pair of equations (4.17).

Proof: From the discussion preceeding Theorem 4.3, it was seen that

$$\mathsf{ERE^{-1}} = \begin{bmatrix} \mathsf{I} & \mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix} \begin{bmatrix} \mathsf{A} & \mathsf{C} \\ \mathsf{O} & \mathsf{B} \end{bmatrix} \begin{bmatrix} \mathsf{I} & -\mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix} = \begin{bmatrix} \mathsf{A} & \mathsf{O} \\ \mathsf{O} & \mathsf{B} \end{bmatrix} = \mathsf{R}'$$

for X satisfying (4.15). By substitution,

$$f_A(ERE^{-1}) = f_A(R')$$
 (4.18)

From Theorem 4.8,

$$f_{\alpha}(R') = \begin{bmatrix} f_{\alpha}(A) & 0 \\ * & f_{\alpha}(B) \end{bmatrix}$$

= $\begin{bmatrix} 0 & 0 \\ * & f_{\alpha}(B) \end{bmatrix}$ (Hamilton-Cayley Theorem 4.9)

From Theorem 4.6

$$f_{A}(ERE^{-1}) = E f_{A}(R) E^{-1} = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix} \begin{bmatrix} U & M \\ V & N \end{bmatrix} \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix}$$

$$= \begin{bmatrix} U+XV & M+XN \\ V & N \end{bmatrix} \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix}$$

$$= \begin{bmatrix} U+XV & -(U+XV)X + (M+XN) \\ V & -VX+N \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ Y & 0 \end{bmatrix} (from (4.18))$$

$$= \begin{bmatrix} 0 & 0 \\ Y & 0 \end{bmatrix} (from (4.18))$$

$$= \begin{bmatrix} 0 & 0 \\ Y & 0 \end{bmatrix} (4.20)$$

Since two matrices are equal iff respective entries are equal, the 1-1th entries of (4.19) and (4.20) imply

$$U + XV = 0 \tag{4.21}$$

and the 1-2th entries imply

$$O = -(U + XV)X + (M + XN)$$

= $OX + (M + XN)$ (from (4.21))
= $M + XN$

Thus a solution of (4.15) will satisfy the pair of equations (4.17).

The example used to illustrate the tensor solution to (4.1) will serve to illustrate Theorem 4.10:

Example Theorem 4.10: Solve the Lypanov equation

$$AX + XB = C$$
 (4.22)
= $AX - X(-B)$

where

A =
$$\begin{bmatrix} s+z & s \\ z & s-z \end{bmatrix}$$
, B = $\begin{bmatrix} -z & -s \\ -z-s & s-z \end{bmatrix}$
C = $\begin{bmatrix} sz & s^2 \\ 2s^2-2z^2-sz & 4sz-2z^2-3s^2 \end{bmatrix}$

From (4.11),

1) .

$$R = \begin{bmatrix} A & C \\ O & -B \end{bmatrix} = \begin{bmatrix} s+z & s & sz & s^2 \\ z & s-z & 2s^2-2z^2-sz & 4sz-2z^2-3s^2 \\ 0 & 0 & z & s \\ 0 & 0 & z+s & z-s \end{bmatrix}$$

Next, find $f_{A}(\mu)$, the characteristic equation of A.

$$f_A(y) = det \begin{bmatrix} s + z - y & s \\ z & s - z - y \end{bmatrix}$$

= $(s+z-y)(s-z-y) - sz$
= $s^2 - sz - sy + zs -z^2 -zy - ys + yz + y^2 - sz$
= $s^2 - sz - z^2 + y^2 - 2sy$

Now evaluate $f_A(R)$ (this was done using the program in Appendix A, a BASIC program performing operations on matrices whose entries are multivariate polynomials):

$$f_{A}(R) = \begin{bmatrix} 0 & 0 & -s^{2}z + 3s^{3} & -5s^{3} + 7s^{2}z - 2sz^{2} \\ 0 & 0 & 5sz^{2} - 5s^{3} - 2z^{3} + 2s^{2}z & 2sz^{2} + 8s^{3} - 8s^{2}z \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 2s^{2} - 2sz & -3s^{2} + 2sz \\ 0 & 0 & -sz - 3s^{2} + 2z^{2} & 5s^{2} - 4sz \end{bmatrix}$$

$$= \begin{bmatrix} 0 & M \\ V & N \end{bmatrix}$$

According to Theorem 4.10, the solution to (4.22) will be found among the common solutions to the pair of equations

$$U + XV = 0$$
 (4.23)
M + XN = 0 (4.24)

Since U = V = 0, (4.23) doesn't contribute to the solution. So the search restricts itself to solutions of the equation

$$M = -XN$$
 or

$$\begin{bmatrix} -s^{2}z + 3s^{3} & -5s^{3} + 7s^{2}z - 2sz^{2} \\ 5sz^{2} - 5s^{3} -2z^{3} + 2s^{2}z & 2sz^{2} + 8s^{3} - 8s^{2}z \end{bmatrix}$$

$$= \begin{bmatrix} w & x \\ y & z \end{bmatrix} \begin{bmatrix} -2s^{2} + 2sz & 3s^{2} - 2sz \\ sz + 3s^{2} - 2z^{2} & -5s^{2} + 4sz \end{bmatrix}$$

However, in order to apply the techniques of Chapter III, one must have an equation of the form

$$AX = b$$

It is an easy matter to convert (4.24) into the form of (3.1) by observing that M = -XN implies $M^T = -N^TX^T$. This latter equation is in the form (3.1), and the techniques of Chapter III can be applied to this transposed equation.

It is possible to further restrict the set of equations within which the solution of (4.15) belongs:

Theorem 4.11: Given the Lypanov equation

$$AX - XB = C A_B_C (c)$$

let R and R' be defined as in (4.11) and $f_{\mbox{\tiny B}}$ the characteristic equation of B. If

$$f_{B}(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

then

$$N - VX = 0$$
 (4.25)
 $M - UX = 0$

Thus a solution to (4.15) will be found among the common

solutions of the pair of equations (4.25).

Proof: The proof is similar to that of Theorem 4.10.

From the discussion preceeding Theorem 4.3, it was seen that

$$\mathsf{ERE^{-1}} = \begin{bmatrix} \mathsf{I} & \mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix} \begin{bmatrix} \mathsf{A} & \mathsf{C} \\ \mathsf{O} & \mathsf{B} \end{bmatrix} \begin{bmatrix} \mathsf{I} & -\mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix} = \begin{bmatrix} \mathsf{A} & \mathsf{O} \\ \mathsf{O} & \mathsf{B} \end{bmatrix} = \mathsf{R}'$$

for X satisfying (4.15). By substitution,

$$f_B(ERE^{-1}) = f_B(R')$$
 (4.26)

From Theorem 4.8,

$$f_B(R') = \begin{bmatrix} f_B(A) & 0 \\ * & f_B(B) \end{bmatrix}$$

$$= \begin{bmatrix} f_B(A) & 0 \\ * & 0 \end{bmatrix}$$
 (Hamilton-Cayley Theorem 4.9) (4.27)

From Theorem 4.6

$$f_{\mathfrak{B}}(\mathsf{ERE^{-1}}) = \mathsf{E} \ f_{\mathfrak{B}}(\mathsf{R}) \ \mathsf{E^{-1}} = \begin{bmatrix} \mathsf{I} & \mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix} \begin{bmatrix} \mathsf{U} & \mathsf{M} \\ \mathsf{V} & \mathsf{N} \end{bmatrix} \begin{bmatrix} \mathsf{I} & -\mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix}$$

$$= \begin{bmatrix} \mathsf{U} + \mathsf{XV} & \mathsf{M} + \mathsf{XN} \\ \mathsf{V} & \mathsf{N} \end{bmatrix} \begin{bmatrix} \mathsf{I} & -\mathsf{X} \\ \mathsf{O} & \mathsf{I} \end{bmatrix}$$

$$= \begin{bmatrix} \mathsf{U} + \mathsf{XV} & -(\mathsf{U} + \mathsf{XV}) \mathsf{X} + (\mathsf{M} + \mathsf{XN}) \\ \mathsf{V} & -\mathsf{VX} + \mathsf{N} \end{bmatrix}$$

$$= \begin{bmatrix} \mathsf{U} + \mathsf{XV} & \mathsf{X} (\mathsf{N} - \mathsf{VX}) + \mathsf{M} - \mathsf{UX} \\ \mathsf{V} & \mathsf{N} - \mathsf{VX} \end{bmatrix} \text{ (rearrange 1-2th entry)}$$

$$= \begin{bmatrix} \mathsf{f}_{\mathfrak{B}}(\mathsf{A}) & \mathsf{O} \\ * & \mathsf{O} \end{bmatrix} \text{ (from (4.27))} \tag{4.29}$$

the 2-2th entries of (4.28) and (4.29) imply

$$N - VX = 0 \tag{4.30}$$

and for the 1-2th entries,

$$O = X(N-VX) + M - UX$$

= $OX + (M - UX)$ (from (4.30))
= $M - UX$

Thus a solution of (4.15) will satisfy the pair of equations (4.25).

The example for Theorem 4.10 will serve to illustrate Theorem 4.11:

Example of Theorem 4.11. From (4.22) begin by finding the characteristic equation for

$$-B = \begin{bmatrix} z & s \\ z+s & z-s \end{bmatrix}$$

$$f_{-B}(y) = det(-B - yI) = det\begin{bmatrix} z-y & s \\ z+s & z-s-y \end{bmatrix}$$

= $(z-u)(z-s-y) - s(z+s)$
= $z^2 - zs - zy - yz + ys + y^2 - sz - s^2$
= $(z^2 - s^2 - 2sz) + (s - 2z)y + y^2$

Using the program in Appendix A, $f_{-8}(R) =$

The pair of equations which need to be satisfied are

$$N - VX = 0$$
 (4.31)
 $M - UX = 0$ (4.32)

(4.31) doesn't contribute to the solution, and so the solution to (4.15) will be found in the solution space of (4.32) M = UX or

$$\begin{bmatrix} -2sz^{2}+2s^{2}z+3s^{3} & 5s^{2}z-2s^{3}-2sz^{2} \\ -5s^{2}z+2z^{3}+s^{3}+sz^{2} & -12sz^{2}+4z^{3}+10s^{2}z-s^{3} \end{bmatrix}$$

$$= \begin{bmatrix} s^{2} & 3s^{2}-2sz \\ -2z^{2}+3sz & 4z^{2}-6sz+s^{2} \end{bmatrix} \begin{bmatrix} w & x \\ y & z \end{bmatrix}$$

To summarize, given the Lypanov equation

$$AX - XB = C A, B, C, X \in \Phi^{\times}(\Gamma)$$

and

$$R = \begin{bmatrix} A & C \\ O & B \end{bmatrix}$$

and $f_A(p)$, $f_B(p)$ the characteristic equations of A and B respectively, then X will be a common solution to the pair of equations

$$M = -XV$$

where

$$f_{A}(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
 and $f_{B}(R) = \begin{bmatrix} U' & M' \\ V' & N' \end{bmatrix}$

Riccati Equation. Attention will now be turned towards (4.2), where $A,B,C,D,X \notin C^{n\times n}(P)$ As with the Lypanov equation, a few insights (not necessarily connected with solving the Riccati equation) will serve to motivate the techniques used to solve (4.2).

If X is an n X n matrix and $I = I_{n \times n}$, then the matrix

$$E = \begin{bmatrix} X & I \\ I & 0 \end{bmatrix} \tag{4.33}$$

always has a determinant equal to -1 (this follows from the permutational definition of the determinant), and is thus unimodular. E's unimodular inverse is

$$E_{\cdot} = \begin{bmatrix} I & -X \\ O & I \end{bmatrix}$$

since EE' =
$$\begin{bmatrix} X & I \\ I & O \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$
 = $\begin{bmatrix} I & O \\ O & I \end{bmatrix}$

and E'E = I likewise follows.

Given the matrix

$$R = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \tag{4.34}$$

where A,B,C,D are nXn matrices,

$$ERE^{-1} = \begin{bmatrix} X & I \\ I & O \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$

$$= \begin{bmatrix} XA+C & XB+D \\ A & B \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$

$$= \begin{bmatrix} XB+D & XA+C-XBX-DX \\ B & A-BX \end{bmatrix}$$

$$= \begin{bmatrix} XB+D & -XBX-DX+XA+C \\ B & A-BX \end{bmatrix}$$
(4.35)

It can be seen that the 1-2th entry of ERE⁻¹ is in the form of the Riccati equation (4.2). However, to get the 1-2th entry to agree exactly with (4.2) which is

$$XDX + AX + XB + C = 0$$

= $-XDX - AX - XB - C$ (4.36)

it will be necessary to modify the R matrix (4.34). The modification is implemented by comparing the form of the 1-2th entry of (4.35)

$$-XBX - DX + XA + C$$
 (from R)

with the the second equation in (4.36)

By matching the coefficient matrices A,B,C,D for like expressions involving X, the suggested change is from

$$R = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \quad \text{to} \quad R = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix}$$
 (4.37)

So with this new form of R,

$$\mathsf{ERE^{-1}} \quad = \quad \begin{bmatrix} \mathsf{X} & \mathsf{I} \\ \mathsf{I} & \mathsf{O} \end{bmatrix} \quad \begin{bmatrix} -\mathsf{B} & \mathsf{D} \\ -\mathsf{C} & \mathsf{A} \end{bmatrix} \quad \begin{bmatrix} \mathsf{O} & \mathsf{I} \\ \mathsf{I} & -\mathsf{X} \end{bmatrix}$$

$$= \begin{bmatrix} -XB-C & XD+A \\ -B & D \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$

$$= \begin{bmatrix} XD+A & -XB-C-XDX-AX \\ D & -B & DX \end{bmatrix}$$

$$(4.38)$$

If X is a solution of the 1-2th entry, then

$$-XB - C - XDX - AX = 0 = XDX + AX + XB + C$$
 and X is a solution of (4.2).

The next insight stems from the procedure followed in solving the Lypanov equation

$$AX - XB = C$$

In Theorems 4.10 and 4.11, a solution to the Lypanov equation was found among the solutions to the pair of equations (4.17)

$$U + XV = 0$$
$$M + XN = 0$$

(derived from using the characteristic equation of A f_A) and (4.25)

$$N' - U'X = 0$$

 $M' - U'X = 0$

(derived from using the characteristic equation of B $f_{\mathfrak{D}}$). Although only one equation from each pair turned out to be significant in solving (4.1), the original pairs point to the solution of the Riccati equation. To obtain (4.17), notice that if

$$\begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} U & M \\ V & N \end{bmatrix} = \begin{bmatrix} U + XV & M + XN \\ V & N \end{bmatrix}$$

is set equal to

then matching the top rows will yield (4.17). The matrix

was picked because it helped generate the Lypanov equation in (4.10). As before,

$$f_A(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
, $R = \begin{bmatrix} A & C \\ O & B \end{bmatrix}$

The development of the pair (4.25) is similar: set

$$\begin{bmatrix} U & M \\ V & N \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix} = \begin{bmatrix} U & -UX + M \\ V & -VX + N \end{bmatrix}$$

equal to

$$\begin{bmatrix} U & O \\ V & O \end{bmatrix} \quad \text{where} \quad f_B(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

to obtain the pair (4.25).

The same approach will now be used to generate solutions to the Riccati equation. Let f(x) be any polynomial in the indeterminate x with degree ≥ 1 . From (4.37)

$$R = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix}$$

Let
$$f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
 (4.39)

Now the crucial step: as if developing the pair (4.17), let X be such that

$$Ef(R) = \begin{bmatrix} X & I \\ I & O \end{bmatrix} \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
$$= \begin{bmatrix} XU+V & XM+N \\ U & M \end{bmatrix} = \begin{bmatrix} O & O \\ U & M \end{bmatrix} = S$$
(4.40)

That such an X exists is motivated by the previous discussion concerning the Lypanov equation. From (4.40) 4.19

$$f(R) = E^{-1}S$$
 or

$$f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix} = \begin{bmatrix} O & I \\ I & -X \end{bmatrix} \begin{bmatrix} O & O \\ U & M \end{bmatrix} = \begin{bmatrix} U & M \\ -XU & -XM \end{bmatrix}$$
(4.41)

Because f(x) is a polynomial in x, xf(x) = f(x)x and so from (4.41)

$$Rf(R) = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix} \begin{bmatrix} U & M \\ -XU & -XM \end{bmatrix} = \begin{bmatrix} -BU-DXU & -BM-DXM \\ -CU-AXU & -CM-AXM \end{bmatrix} (4.42)$$

and

$$f(R)R = \begin{bmatrix} U & M \\ -XU & -XM \end{bmatrix} \begin{bmatrix} -B & D \\ -C & A \end{bmatrix} = \begin{bmatrix} -UB-MC & UD+MA \\ XUB+XMC & -XUD-XMA \end{bmatrix}$$
(4.43)

Since (4.42) and (4.43) are the same matrix

$$BU + DXU = UB + MC$$
 (4.44)
 $BM + DXM = -UD - MA$ (4.45)
 $CU + AXU = -XUB - XMC$ (4.46)
 $CM + AXM = XUD + XMA$ (4.47)

Within the above four equations lie two solutions to the Riccati equation (4.2). From (4.44) and (4.46),

CU + AXU = -X(UB + MC) = -X(BU + DXU) = -XBU - XDXUand so

$$CU + AXU + XBU + XDXU = 0$$

$$= (C + AX + XB + XDX)U \qquad (4.48)$$

If det(U) is a nonzero complex number, then U^{-1} exists and from (4.48)

$$XDX + AX + XB + C = O$$

and X satisfies the Riccati equation (4.2). The second solution uses (4.45) and (4.47):

CM + AXM = X(UD + MA) = X(-BM - DXM) = -XBM - XDXMand so

$$CM + AXM + XBM + XDXM = O = (C + AX + XB + XDX)M$$

If det(M) is a nonzero complex number, M^{-1} exists and so XDX + AX + XB + C = 0

another solution to (4.2).

The following theorem due to Jones has now been proven:

Theorem 4.12: Given A,B,C,D ($\phi^{n\times n}(P)$ and f(x) a polynomial with degree ≥ 1 , let

$$R = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix} \qquad \text{and} \qquad f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

If $X \in \Phi^{n \times n}(\Gamma)$ satisfies

$$XU + V = 0$$
 (4.49)
 $XM + N = 0$

and det(M) $\{ (-10) \}$, then XDX + AX + XB + C = 0. Also, if X $\{ (-10) \}$ satisfies the pair (4.49) and det(U) $\{ (-10) \}$ then XDX + AX + XB + C = 0.

Proof: previous discussion.

A third solution exists for (4.2) but requires a different setup akin to developing the pair (4.25). Let R be as in (4.37) and f(R) as in (4.39). Let X satisfy

$$f(R) E = \begin{bmatrix} U & M \\ V & N \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix} = \begin{bmatrix} M & U-MX \\ N & V-NX \end{bmatrix}$$
$$= \begin{bmatrix} M & O \\ N & O \end{bmatrix}$$
(4.50)

That such an X exists is motivated by the development of the pair (4.25) for the Lypanov equation. (4.50) implies

$$f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix} = \begin{bmatrix} M & O \\ N & O \end{bmatrix} \begin{bmatrix} X & I \\ I & O \end{bmatrix} = \begin{bmatrix} MX & M \\ NX & N \end{bmatrix}$$
 (4.51)

Since Rf(R) = f(R)R

$$Rf(R) = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix} \begin{bmatrix} MX & M \\ NX & N \end{bmatrix}$$
$$= \begin{bmatrix} -BMX + DNX & -BM + DN \\ -CMX + ANX & -CM + AN \end{bmatrix}$$
(4.52)

and

$$f(R) R = \begin{bmatrix} MX & M \\ NX & N \end{bmatrix} \begin{bmatrix} -B & D \\ -C & A \end{bmatrix}$$
$$= \begin{bmatrix} -MXB - MC & MXD + MA \\ -NXB - NC & NXD + NA \end{bmatrix}$$
(4.53)

Equating last rows of (4.52) and (4.53)

$$NXB + NC = CMX - ANX$$

$$NXD + NA = -CM + AN$$

$$(4.54)$$

$$(4.55)$$

It follows that

$$NXB + NC = (CM - AN)X = (-NXD - NA)X$$

= $-NXDX - NAX$

and so

$$NXDX + NAX + NXB + NC = 0 = N(XDX + AX + XB + C)$$

If det(N) is a nonzero complex number, N^{-1} exists and so

$$XDX + AX + XB + C = 0$$

and a third solution to (4.2) has been found and thus

$$R = \begin{bmatrix} -B & D \\ -C & A \end{bmatrix} \qquad \text{and} \qquad f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

If $X \in C^{n \times n}(\Gamma)$ satisfies

$$U - MX = 0$$
 (4.56)
 $V - NX = 0$

and det(N) $f = \{0\}$, then XDX + AX + XB + C = 0.

Proof: previous discussion.

The next theorem presents a companion Riccati equation

to that of (4.2) given a particular polynomial f(x):

Theorem 4.14: Let f(x) be a polynomial of degree ≥ 1 and

$$XDX + AX + XB + C = O$$
 A,B,C,D,X $\notin \Phi^{n\times n}(P)$

with R and f(R) given by (4.37) and (4.39). Then

$$XMX + NX - XU - V = 0$$

Proof: It has already been shown that

$$E = \begin{bmatrix} X & I \\ I & 0 \end{bmatrix} \text{ and } E' = \begin{bmatrix} 0 & I \\ I & -X \end{bmatrix}$$

are both unimodular and inverses of one another. From (4.38) and the given,

$$ERE^{-1} = \begin{bmatrix} XD + A & O \\ D & -B - DX \end{bmatrix}$$

From Theorem 4.8,

$$f(ERE^{-1}) = \begin{bmatrix} f(XD+A) & O \\ * & f(-B-DX) \end{bmatrix}$$

$$= Ef(R)E^{-1} \quad (from Theorem 4.6)$$

$$= \begin{bmatrix} X & I \\ I & O \end{bmatrix} \begin{bmatrix} U & M \\ V & N \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$

$$= \begin{bmatrix} XU + V & XM + N \\ U & M \end{bmatrix} \begin{bmatrix} O & I \\ I & -X \end{bmatrix}$$

$$= \begin{bmatrix} XM + N & XU + V - XMX - NX \\ M & U - MX \end{bmatrix} \quad (4.58)$$

Equating the 1-2th entries of (4.57) and (4.58),

$$XU + V - XMX - NX = O = XMX + NX - XU - V$$

V Solutions of the Third Order Riccati Equation XAXBX + XCX + XD + EX + F = 0

This chapter will discuss some of the approaches which may yield solutions to the third order Riccati equation

$$XAXBX + XCX + XD + EX + F = 0$$
 (5.1)

where A,B,C,D,E,F,X $\notin \Phi^{n\times n}(\Gamma)$, Γ the tuple of variates. The approach for (5.1) will be less general than that for the second order Riccati and Lypanov equations because of the form of (5.1). Difficulties stemming from this form are discussed in the next chapter.

One approach to solving (5.1) is akin to that taken for the second order Riccati. Let A,B,C,D,E,F be known matrices in $\P^{n\times n}(\mathbb{P})$ and

$$R = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$$

and

$$f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

where f(x) is a given polynomial with degree ≥ 1 . As seen previously, if $X \in \Phi^{\times n}(\Gamma)$,

$$\begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} U & M \\ V & N \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix} = \begin{bmatrix} U + XV & -(U + XV)X + (XN + M) \\ N - VX \end{bmatrix}$$

Suppose U + XV = 0 (motivated by its appearance in solving for the Lypanov and second order Riccati equations). Thus

$$U = -XV \tag{5.2}$$

Because f(R) is a polynomial in R, R and f(R) commute. From (5.2),

$$f(R) = \begin{bmatrix} -XV & M \\ V & N \end{bmatrix}$$

It follows that

$$Rf(R) = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} -XV & M \\ V & N \end{bmatrix} = \begin{bmatrix} BV - AXV & AM + BN \\ DV - CXV & CM + DN \end{bmatrix}$$

and

$$f(R)R = \begin{bmatrix} -XV & M \\ V & N \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} MC - XVA & MD - XVB \\ VA + NC & VB + ND \end{bmatrix}$$

Equating the first columns,

$$BV - AXV = MC - XVA$$
 (5.3)

$$DV - CXV = VA + NC$$
 (5.4)

VA appears in the right hand sides of (5.3) and (5.4). If MC in (5.3) involved XNC, then a substitution from (5.4) could be made in (5.3). Accordingly, let

$$MC = -XNC - \Sigma \tag{5.5}$$

Then (5.3) becomes

BV - AXV =
$$(-XNC - \Sigma) - XVA$$

= $-X(NC + VA) - \Sigma$
= $-X(DV - CXV) - \Sigma$ (from (5.4))
= $-XDV + XCXV - \Sigma$

or

$$XCXV - XDV + AXV - BV - \Sigma = 0$$
 (5.6)

Because all terms except E end in V, let

$$\Sigma = e_1 V$$

(5.6) then becomes

$$(XCX - XD + AX - B - e_1)V = 0$$

If det(V) is a complex, non-zero number (easily verified since f(R) is known), then the inverse of V exists and so

$$XCX - XD + AX - B - e_1 = 0$$
 (5.7)

Since solutions of the third order Riccati equation are

sought, let $e_1 = -XEXFX$. (5.7) then becomes

$$XEXFX + XCX + AX - XD - B = 0 (5.8)$$

and X is a solution to this third order Riccati equation.

However, since $e_1=-XEXFX$ and $\Sigma=e_1V$, it follows that $\Sigma=-XEXFXV$ and from (5.2), $\Sigma=XEXFU$. From (5.5) it follows that

$$MC = -XNC - \Sigma = -XNC - XEXFU$$

To summarize, if

$$R = \begin{bmatrix} A & B \\ C & D \end{bmatrix} , \qquad f(R) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$

det(V) a non-zero complex number and X satisfies the pair of
equations

$$U + XV = 0$$

$$-XEXFU - XNC = MC$$
 (5.9)

then X is a solution to the third order Riccati equation

$$XEXFX + XCX + AX - XD - B = 0 (5.10)$$

If det(FU) is a non-zero complex number, then (5.9) can be reduced to a second order Riccati equation of the form solved in the previous chapter.

Another solution to (5.10) can be gleened using the above approach. Let XN + M = 0. Then

$$Rf(R) = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} U & -XN \end{bmatrix} = \begin{bmatrix} AU + BV & BN - AXN \\ CU + DV & DN - CXN \end{bmatrix}$$
(5.11)

and

$$f(R)R = \begin{bmatrix} U & -XN \\ V & N \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} UA - XNC & UB - XND \\ VA + NC & VB + ND \end{bmatrix} (5.12)$$

Equating the second columns of (5.11) and (5.12),

$$-AXN + BN = UB - XND$$
 (5.13)
 $-CXN + DN = VB + ND$ (5.14)
5.3

Noticing that ND appears in the right hand sides, let

$$UB = -XVB + \Sigma \tag{5.15}$$

Then (5.13) becomes

$$-AXN + BN = (-XVB + \Sigma) - XND$$

$$= -X(VB + ND) + \Sigma$$

$$= -X(DN - CXN) + \Sigma$$

$$= -XDN + XCXN + \Sigma$$
(5.16)

and so

$$XCXN - XDN + AXN - BN + \Sigma = 0$$
 (5.17)

Let $\Sigma = e_1N$. Then (5.17) becomes

$$(XCX - XD + AX - B + e_1)N = 0$$

If det(N) is a non-zero complex number, then N has an inverse, and so

$$XCX - XD + AX - B + e_1 = 0$$
 (5.18)

If e₁ = XEXFX then

$$XCX - XD + AX - B + XEXFX = 0$$

and another solution to the third order Riccati has been found. Since $\Sigma = e_1N$, $\Sigma = XEXFXN = -XEXFM$. (5.15) then becomes

Thus, if X is a common solution to the pair of equations

and det(N) is non-zero complex, then

$$XEXFX + XCX + AX - XD - B = O$$

Another approach to solving (5.1) extends the solution via similarity transformation used in the matrix equations presented herein. Though this approach fails to generate 5.4

sets of equations (whose common solutions satisfy (5.1)), it does present relations among the coefficient matrices for a restricted case of (5.1).

Let A,B,C,D,E,F,G,H ($^{n\times n}(P)$. Then given X ($^{n\times n}(P)$.

$$\begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} E & F \\ G & H \end{bmatrix} \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix}$$

$$= \begin{bmatrix} E+XG & F+XH \\ G & H \end{bmatrix} \begin{bmatrix} A+XC & -AX-XCX+B+XD \\ C & D-CX \end{bmatrix}$$

= 5

By multiplying these matrices, the following entries of are obtained:

1-1: EA + EXC + XGA + XGXC + FC + XHC (5.19)
1-2:
$$-XGXCX - X(GA+HC)X - (EA+FC)X + X(GB+HD) +$$

$$1-3: GA + GXC + HC$$
 (5.21)

Of the entries, the only one in which the "cubic" of the third order Riccati appears is (5.20), that is, XGXCX. If it weren't for the last term in paranthesis in (5.20), the form of (5.20) would be the same as that of (5.1). In order to eliminate this last term and preserve the structural integrity of (5.20), let D = E = 0. It follows that

$$S = \begin{bmatrix} XGA+XGXC+FC+XHC & -XGXCX-X(GA+HC)X-FCX+XGB \\ GA+GXC+HC & GB-GAX-GXCX-HCX \end{bmatrix}$$
 (5.22)

If X were a solution to $\S(1,2)$, then X would be a solution to a third order Riccati equation of the form

$$XAXBX + XCX + XD + EX = 0 (5.23)$$

which is a special case of (5.1). From the remarks preceding (5.19), it follows that (since D = E = 0)

$$\xi = \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} O & F \\ G & H \end{bmatrix} \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} A & B \\ C & O \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix}$$
(5.24)

$$= \begin{bmatrix} XGA+XGXC+FC+XHC & 0 \\ GA+GXC+HC & GB-GAX-GXCX-HCX \end{bmatrix}$$
 (5.25)

Multiplying the three middle matrices in (5.24),

$$\begin{bmatrix} O & F \\ G & H \end{bmatrix} \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} A & B \\ C & O \end{bmatrix} = \begin{bmatrix} FC \\ GA + HC + GXC & GB \end{bmatrix}$$

$$= \overline{g}_1$$

If f(x) is any polynomial of degree 2 1, then from (5.24) and Theorem 4.6,

$$f(\delta) = \begin{bmatrix} I & X \\ O & I \end{bmatrix}$$
 $f(\delta_1)$ $\begin{bmatrix} I & -X \\ O & I \end{bmatrix}$

and from Theorem 4.8

$$f(\xi) = \begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} f(FC) & O \\ * & f(GB) \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix}$$

$$= \begin{bmatrix} f(XGA+XGXC+FC+XHC) & O \\ * & f(GB-GAX-GXCX-HCX) \end{bmatrix}$$

Since $f(\S)$ and $f(\S_1)$ are similar matrices, they have the same determinants. From the permutational definition of the determinant (2.13),

and

$$det(f(\S_1)) = det[f(FC)] * det[f(GB)]$$
 (5.29)

and thus

The following theorem has now been proven:

Theorem 5.1: Given A,B,C,F,G,H ($\Phi^{n\times n}(P)$ and X ($\Phi^{n\times n}(P)$) a solution to

-XGXCX-X(GA+HC)X-FCX+XGB = 0

then

Proof: previous discussion.

Theorem 5.1 may be extended to a Riccati equation of the form

$$XA_1XA_2X + XA_3X + A_4X + XA_5 = 0$$
 (5.30)

To get this equation in the form (of the given in Theorem 5.1)

-XGXCX-X(GA+HC)X-FCX+XGB = 0

equating coefficient matrices is done. Thus

$$A_1 = -G$$
 (5.31)
 $A_2 = C$ (5.32)
 $A_3 = -(GA + HC) = A_1A - HA_2$ (5.33)
 $A_4 = -FC = -FA_2$ (5.34)
 $A_5 = GB = -A_1B$ (5.35)

Since G and C are readily known, it remains to determine A,H,F and B. From the above equations, one may have considerable latitude in choosing A,H,F,B. From (5.35), B satisfies the equation $A_1B = -A_5$. Likewise, F satisfies the equation $FA_2 = -A_4$. Lastly, A and H satisfy the Lypanov equation (5.33)

By proper choice of A,B,F,H, the challenge of finding solutions to (5.30) via Theorem 5.1 may be made easier.

VI Conclusions and Recommendations

This paper sought the solution X to four types of matrix equations: the linear equation

$$AX = b$$

the Lypanov equation

$$AX - XB = C$$

the second-order Riccati equation

$$XDX + AX + XB + C = 0$$

and the third-order Riccati equation

$$XAXBX + XCX + XD + EX + F = 0$$

where the entries of all matrices are restricted to being multivariate polynomials over the complex numbers. The method of solution involved two phases:

- identify a similarity transformation on a matrix Σ in which is embedded the coefficient matrices of the above equations. The transformation gives a matrix whose entries include the equation being solved.
- 2. identify a polynomial which, when its argument is the matrix Z in (1), gives a matrix whose entries yield pairs of linear equations. The common solutions of these pairs of equations will contain those of the given matrix equation. The choice of polynomial may yield different solutions.

Since the search for solutions ends in solving a linear equation, a method was presented which may identify the general solution of the given linear equation. The method's success depends on if the matrix A in AX = b is reducible to a Smith Form. The similarity transformation of (1) is given by E Σ E⁻¹ where E equals

$$\begin{bmatrix} 0 & 1 \end{bmatrix}$$

(which is unimodular) and ${f \Sigma}$ for the Lypanov equation equals

and Σ for the second order Riccati equals

The third order Riccati presented difficulties because its form wasn't adaptable to the method of similarity transformations. Nevertheless, a transformation could be done which might yield some solutions. When successful, the solution to the third order Riccati was a common solution to a linear and a second order Riccati equation. The polynomial in (2) can assume any form and when its argument is Σ ,

$$f(\Sigma) = \begin{bmatrix} U & M \\ V & N \end{bmatrix}$$
 (6.1)

The submatrices U,M,V,N yield pairs of equations which depend on the matrix equation being solved. These pairs in turn yield solutions of the original matrix equation.

Recommendations. The biggest challenge is also the central concern: develop the mathematical theory which will identify whether or not a given matrix, whose entries are multivariate polynomials, is reducible to Smith Form.

Though the answer for the two-variate polynomial case is known, the three or more variate case remains an open question. It's felt that this challenge will involve

creative work involving number theory, abstract algebra, and matrix theory. Once the theory identifying the conditions under which reducibility to Smith Form is known, the next hurdle will be to develop a computer algorithm performing the reductions (the Smith Form may not be unique). The papers [8] and [20] give approaches to the two-variate case, and the papers [23] and [33] give insights dealing with the "computer programming algebra" of matrix forms and polynomials. Perhaps there is an underlying linear programming or network formulation to the Smith Form reduction.

In solving the Lypanov and Riccati equations, it was necessary to assume the existence of the inverse to one or more of the submatrices in (6.1). Of course, (6.1) depends on the polynomial f(x) chosen, which in turn dictates the types of solutions found to the original matrix equation. Research needs to be done into the types of polynomials yielding one or more invertible submatrices of (6.1). Perhaps there are equivalence classes of polynomials. Or families of polynomials. Maybe there is no structure. Or perhaps there is a "minimum" polynomial which (in some way) generates all polynomials yielding invertible submatrices. Again, a knowledge of abstract algebra may prove valuable: much literature dealing with polynomial structures is couched in abstract algebraic terms.

Much effort was spent in finding the "proper embedding"

for the third order Riccati equation. Disappointingly limited success was realized. To see what is meant by "proper embedding", notice that in

$$\begin{bmatrix} I & X \\ O & I \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I & -X \\ O & I \end{bmatrix} = \begin{bmatrix} A + XC & -AX - XCX + B + XD \\ C & -CX + D \end{bmatrix}$$

a second order Riccati equation is found in the 1-2th position of the matrix on the right. If C = 0, then this equation becomes the Lypanov. Thus the Lypanov and second order Riccati equations are "embedded" in the matrix on the right, which in turn comes from the similarity transformation on the left. Since this approach has proven so successful in generating linear equations whose solutions are those to the Lypanov and second order Riccati equations, it is natural to extend the similarity transformation to handle the third order Riccati equation.

Unfortunately, this is more easily said than done. Since the third order Riccati equation

$$XAXBX + XCX + XD + EX + F = 0$$
 (6.2)

has six coefficient matrices, the above approach would have to be modified to handle these six. As it stands, it is geared to the four coefficients of the second order Riccati equation. Is it possible to work with matrices which are two sub-matrices deep and wide, or is it necessary to go to higher order matrices, perhaps three submatrices in dimension. If so, what would the unimodular matrix be that effects the similarity transformation? Is the restriction of unimodularity unduly restrictive, mandatory though it

seems? Also, the first term XAXBX of (6.2) presents a challenge: how to introduce the middle X from the transformation E Σ E⁻¹ while hopefully embedding a third order Riccati equation in the resulting matrix of transition. Perhaps a different type of transformation needs to be done. Instead of a single transform, a string of transforms may be the answer, for example

E121E222E3

where the \$\mathbb{Z}_{\text{\tinx{\text{\til\text{\te

Or should it? If not embedded, could other matrix equations which are embedded as a matter of course yield, in some combination, some (or all!) solutions of the third order Riccati? Another concern arises: in the approaches presented herein, a specially selected polynomial f(x) played the crucial role of generating pairs of equations whose common solutions included those of the matrix equation in question. For the third order Riccati, how is the polynomial to be selected? How should the pairs be generated, if indeed this pattern holds? Perhaps triplets of equations arise instead of pairs.

The unsettling thought occurs: maybe similarity transforms are not powerful enough for the third order

Riccati, and whole new approaches are in order. After all, current research into these matrix equations is still in the pioneering stages. Hopefully, the successful resolution to the third order Riccati equation will point the way to solving higher order Riccati equations. However, a spectre appears: in solving for the roots of a polynomial in one variable, the insolvability of the quintic was demonstrated by Abel in the early nineteenth century. Could there be a similar obstacle ahead for matrix equations? In any case, finding solutions to matrix equations will prove challenging and endlessly fascinating—as well as having immediate practical applications.

The last recommendation is a minor one: create a new segment (in the computer program of Appendix A) which will compute the determinant of a given square matrix having multivariate polynomial entries. The method would have to keep calculations down to a minimum, since finding determinants is computationally intensive. One approach, which seems to be the quickest way to find a matrix's determinant, was published by G. Macloskie [22] in 1904. This method could be adapted to that of finding determinants for matrices with polynomial entries.

Appendix A: BASIC program performing matrix operations

The following BASIC program performs operations on rectangular matrices having multivariate polynomial entries over the real numbers. Via menu options, the interactive program allows the user to

- -create a matrix
- -view a matrix
- -transpose a matrix
- -add two matrices
- -multiply two matrices
- -extract the U,M,V,N matrices from a given matrix

 The size of matrices which the program can handle is limited only by the amount of memory available on the computer and the parameters in the DIMENSION statement. The program was designed on an IBM PC (DOS 3.0) and was intended to assist the thesis effort.

The program's logic hinges upon the way it recognizes a polynomial in n-variates: as a set of (n+1)-tuples each of whose entries come from the coefficients and exponents of the polynomial. To obtain the tuples, the entire polynomial is re-expressed in "standard form" as a sum of terms, each having all variate and exponent positions appear. For instance, the polynomial in the variates x and y

$$2x^2y - 17xy^2 - 3y + 7$$
 (A.1)

is re-expressed in standard form as

$$2x^2y^1 + (-17x^1y^3) + (-3x^0y^1) + (7x^0y^0) \qquad (A.2)$$
 and is the sum of the four terms $2x^2y^1$, $-17x^1y^3$, $-3x^0y^1$,
$$7x^0y^0$$
. Though the order of the variates (i.e., x before y

versus y before x) is arbitrary, the order must be consistent.

Once in standard form, each term is put into its tuple representation

(coefficient, exponent,...,exponent,) (A.3) with the polynomial represented by the collection of the tuples. For the above example:

TERM	3-tuple
	
2x2y1	(2,2,1)
-17x 1y=	(-17,1,5)
-3x°y1	(-3,0,1)
7x°y°	(7,0,0)

and so the polynomial

$$2x^2y - 17xy^3 - 3y + 7$$

is represented by the set

$$\{(2,2,1), (-17,1,5), (-3,0,1), (7,0,0)\}$$

Similarly, the polynomial

is re-expressed in the standard form (variate order is x,y,z,u,v)

$$42x^{1}y^{1}z^{1}u^{1}v^{1} + (-2x^{1}y^{1}z^{2}u^{0}v^{0})$$

and thus the set of 6-tuples

Addition and multiplication between two polynomials are easily expressed in terms of their tuple representations provided

-all tuples have the same number of entries
-coordinate positions all correspond to the same
variable

Addition. Given two terms in n-variates, let (a, α) and (b, β) be their respective (n+1)-tuple representations where a, b are the coefficients of each term and α, β the exponent vectors. For example, if the terms are $3x^2y$ and $7x^2y^2$, then

$$3x^2y \approx (3,2,1)$$
, $a = 3$, $\alpha = (2,1)$
 $7x^2y^2 \approx (7,9,5)$, $b = 7$, $\beta = (9,5)$

Tuple addition is defined by

The operation # is clearly commutative. To add two polynomials, the tuples representing the terms are added according to (A.4). For example:

$$(2x^2y - 9xy^2) + (4xy^2 + x^2y)$$

% [(2,2,1) # (-9,1,2)] # [(4,1,2) # (1,2,1)]

then, rearranging terms

=
$$[(2,2,1) \div (1,2,1)] \div [(-9,1,2) \div (4,1,2)]$$

= $(3,2,1) \div (-5,1,2)$
 $(3,2,1), (-5,1,2)$
 $3x^2y - 5xy^2$

<u>Multiplication</u>. Multiplication between two tuples is defined by

$$(\mathbf{a}, \mathbf{C}) \bullet (\mathbf{b}, \mathbf{B}) = (\mathbf{a}\mathbf{b}, \mathbf{C} + \mathbf{B}) \tag{A.5}$$

where $\alpha + \beta$ is the familiar operation of addition between two vectors in Euclidean n-space. For example,

$$(2x^{3}y^{5})(6x^{7}y^{6}) \approx (2,3,5) \cdot (6,7,8)$$

= $(2*6,3+7,5+8)$
= $(12,10,13)$
 $\approx 12x^{10}y^{13}$

Using (A.5) the tuple representation of the product between two polynomials is quickly found. Consider the product of the n-term polynomial

$$TERM_{11} + \dots + TERM_{10} \tag{A.6}$$

and the m-term polynomial

$$TERM_{21} + \dots + TERM_{2m} \tag{A.7}$$

which is

If the value of TERM_{Jk} is now changed to the tuple form of the polynomial element TERM_{Jk}, then the product of (A.6) and (A.7) expressed in tuple form is

EXAMPLE:

$$\begin{array}{l} (x^2y^3 + 12x^4y^3) (x^4y^7 + 11x^3y^7) \\ &= (x^2y^3) (x^4y^7) + (x^2y^3) (11x^3y^7) \\ &+ (12x^4y^3) (x^4y^7) + (12x^4y^3) (11x^3y^7) \\ &\approx \left[(1,2,3) \cdot (1,6,7) \right] \, \div \left[(1,2,3) \cdot (11,8,9) \right] \\ &+ \left[(12,4,5) \cdot (1,6,7) \right] \, \div \left[(12,4,5) \cdot (11,8,9) \right] \\ &= (1,8,10) \, \div \left(11,10,12 \right) \\ &+ \left(12,10,12 \right) \, \div \left(132,12,14 \right) \\ &= \left(1,8,10 \right) \, \div \left(23,10,12 \right) \, \div \left(132,12,14 \right) \\ &= \left(1,8,10 \right) \, \cdot \left(23,10,12 \right) \, \cdot \left(132,12,14 \right) \\ &\approx x^3y^{10} + 23x^{10}y^{12} + 132x^{12}y^{14} \end{array}$$

Network. Put in a nutshell, the program keeps track of tuple operations using a system of pointers (program lines 1150-1490). That is, a network tree is constructed which A.4

represents the results of tuple operations. This section assumes knowledge of networks as given in [10:91-124].

The fundamental insight is that a polynomial can be represented by a tree. Specifically, the results of tuple operations is maintained in a network tree whose nodes represent a particular variate in a polynomial term. Nodal "potentials" are the exponent of the corresponding variate. Terminal nodes of the polynomial's tree have a second potential whose value is that of the coefficient of a particular term. The i'th depth of the tree corresponds to the i'th variate. For example, the polynomial

is represented by the tree in Figure 1. All the nodes at depth one are x variates; nodes at depth two are y variates; nodes at depth three are z variates. The numbers in [] are node potentials, and numbers in []* are the coefficient of the given polynomial term. From this example, it can be seen that a polynomial term corresponds to a unique path from node 0 to a given terminal node at depth three, e.g., the term $8x^{12}y^{7}z^{22}$ has the nodal path 0-4-5-7.

In performing a tuple operation, the order in which the branches are built are term by term. The following example illustrates the sequence in which the program builds branches.

EXAMPLE: Construct the tree representing

$$x^{2}(1 + y) + y(x^{2} + y)$$

$$x^{2}(1 + y) + y(x^{2} + y) = x^{2} + x^{2}y + yx^{2} + y^{2}$$

= $x^{2}y^{0} + x^{2}y^{1} + y^{1}x^{2} + x^{0}y^{2}$ (A.9)

Since there are 2 variates x,y the tree will have 2 depths. The first term is x^2y° which contributes the first branch 0-1-2 in Figure 2. The second term x^2y^1 contributes the branch 0-1-3 in Figure 3. Note that the second potential of node 3 ([]*) is 1: the coefficient of the second term x^2y^1 . Because the third term of (A.9) is also x^2y^1 , it doesn't contribute a branch to the tree. However, the second potential at node 3 is increased by 1: the coefficient in the third term x^2y^1 (Figure 4). The fourth term x^0y^2 contributes the branch 0-4-5 in Figure 5. Since there are no more terms in (A.9), the tree in Figure 5 represents the polynomial $x^2(1+y) + y(x^2+y)$


Manual input of data. When creating a matrix via the menu prompt to "huild a matrix", polynomials are entered term by term in their tuple form with no commas between tuple entries. Each tuple MUST remain on the same line, and each tuple except the last ends with a carriage return; the last tuple ends with a semicolon followed by a carriage return. The program will then prompt for the next polynomial entry and the above protocol is repeated. A sample session inputing the matrix


$$Q = \begin{bmatrix} x^2 + 3xyz + z^2 & 4 + 2x - 3y \\ z^2 + 17y^{90} & x^2y^3z^4 \end{bmatrix}$$
 (A.10)

is given in Figure 6.

Output. The program outputs a polynomial in its tuple

form. Figure 7 is a sample session which printed the matrix fA(R) preceding (4.23).





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - 1




Figure 1. Network tree representing the polynomial $3x^2yz^{13} - 7x^{12}y^9z + 8x^{12}y^9z^{22}$




Fig. 2. Branch for x^2




Fig. 3. Branch for $x^2 + x^2y$




Fig. 4. Branch for $x^2 - 2x^2y$




Fig. 5. Branch for $x^2 + 2x^2y + y^2$

```
*****************
MENU:
         Build a new matrix"
         View a matrix"
         Transpose a matrix"
       + Add 2 matrices"
         Multiply 2 matrices"
       U Extract U,M,V,N matrices
? B
Matrix name? B:Q
ROW, COL dimension? 2,2
Number of polynomial variates? 3
Build a square diagonal matrix whose diagonal entries
  are equal Y/N? N
*** COEF EXP1 EXP2 ... EXPn ***
 a semicolon; represents end of polynomial
(1,1):
? 1 2 0 0
? 3 1 1 1
? 1 0 0 2;
(1,2):
? 4 0 0 0
? 2 1 0 0
? -3 0 1 0:
(2,1):
71006
? 17 0 98 0;
(2,2):
? 1 2 3 4;
```

Figure 6. Session building matrix (A.10)

```
********************
MENU:
       B Build a new matrix"
         View a matrix"
         Transpose a matrix"
       + Add 2 matrices"
       * Multiply 2 matrices"
       U Extract U,M,V,N matrices
************
Screen output Y/N? Y
Matrix to view? B:fA(R)
MATRIX B: fA(R) IS 4 X 4 and has 2 polynomial variates
B:fA(R)(1,1):
                               B: fA(R) ( 3, 1):
0 0 0
                               0 0 0
                               B: fA(R) ( 3, 2):
B:fA(R)(1,2):
0 0 0
                               0 0 0
                               B: fA(R) ( 3, 3):
B:fA(R)(1,3):
                               2 2 0
-2 1 1
-1 2 1
3 3 0
                               B: fA(R) ( 3, 4):
B:fA(R)(1,4):
-5 3 0
                               -3 2 0
                                  1
                                2
 7
   2
     1
-2 1
     2
B:fA(R)( 2, 1):
                               B:fA(R)(4,1):
                               0 0 0
0 0 0
B:fA(R)(2, 2):
                               B:fA(R)(4,2):
0 0 0
                               0 0 0
B.fA(R)( 2, 3):
                               B: fA(R) ( 4, 3):
                               -1 1
   1
     2
                                     1
-5
   3 0
                               -3 2 0
-2
                                2 0
                                     2
   0
     3
 2 2 1
B:fA(R)( 2, 4):
                               B: fA(R) ( 4, 4):
                                5 2 0
   1
      2
                                  1
   3
                               -4
 8
     0
   2
-8
```

Figure 7. Session viewing the matrix fA(R) preceeding (4.23)

NOTE: the program has been modified to fit within the format requirements of the AFIT Style Guide, and so doesn't represent the correct syntax of the program. However, the adjustments are few and are apparent.

•)

```
10
        'CPT Bruce W. Colletti, AFIT/EN; 1 August 1985
20
        'The input file for this program (which multiplies 2
30
        'matrices having multivariate polynomial entries)
        'has the following
40
        'structure:
                     row, column dimension of matrix A;
50
            line 1:
                      number of variates
60
            line 2:
                      number of terms in the A(1.1)
                      polynomial
70
            line 3:
                      tuple corresponding to the 1st term of
                      A(1,1)
                      tuple corresponding to the 2nd term of
80
            line 4:
                      A(1,1)
90
            line n:
                     tuple corresponding to the last term
                      of A(1,1)
100
          line n+1:
                     number of terms in the A(1,2)
                      polynomial
110
          line n+2:
                      tuple corresponding to the 1st term of
                      A(1,2)
120
          line n+3:
                     tuple corresponding to the 2nd term of
                      A(1,2)
            line m:
130
                     tuple corresponding to the last term
                      of A(1,2)
140
          line m+1:
                      number of terms in the A(1,3)
                      polynomial
150
               etc:
                      matrix entries are entered row at a
                      time
160
        CLS : CLEAR
170
        ROWS = 5 : COLS = 5 : TERM = 25 : VAR = 5
180
        DIM A(ROWS, COLS, TERM, VAR), B(ROWS, COLS, TERM, VAR),
        B$ (20)
190
        DIM F(90),PDT(90),R(90),BACK(90),M(90),E(VAR)
200
        DIM COEFL (ROWS, COLS), COEFR (ROWS, COLS)
210
        'A contains the elements of the left matrix and B
        'the elements of
220
        'the right (e.g., A + B, A*B):
230
        'A(I,J,K,L) the exponent of the L'th variate of the
        'K'th term of the
240
                     I-J'th polynomial entry of A
250
        'B(I,J,K,L) the exponent of the L'th variate of the
        'K'th term of the
260
                     I-J'th polynomial entry of B
270
            BACK(i) the node-back pointer for node i
280
        'COEFL(i,j) the number of terms in the polynomial
        'located at
290
                     A(i,j)
300
        'COEFR(i,j) the number of terms in the polynomial
        'located at
```

```
310
                    B(i,j)
320
            E(i)
                    contains the exponent of the i'th
                    variable of the
                    product between 2 polynomial terms
330
340
            F(i)
                    forward pointer for node i
350
            M(i)
                    the "summed coefficient". This is a
                    node in the final
360
                    tree with no forward pointer.
                    the coefficient of
370
                    that term of the polynomial having the
                    path from given
380
                    terminal node to source node
390
            POT(i)
                    potential of node i; that is, the
                    exponent of the
400
                    variable represented by the node
410
                    the right pointer for node i
420
        PRINT "************************
430
        PRINT "MENU:"
440
        PRINT "
                        Build a new matrix"
                      В
        PRINT "
450
                         View a matrix"
        PRINT "
460
                         Transpose a matrix"
        PRINT "
470
                         Add 2 matrices"
        PRINT "
480
                         Multiply 2 matrices"
490
        PRINT "
                      U Extract U.M.V.N matrices
        PRINT "***********
500
510
        INPUT OPT$
        IF OPT$ = "B" THEN 2410
520
        IF OPT$ = "V" THEN 2790
530
        IF OPT$ = "U" THEN 3470
540
        IF OPT$ = "'" THEN 3130
550
        IF ((OPT$ <> "*") AND (OPT$ <> "+")) THEN 420
560
570
        INPUT "Left matrix"; FILEL$
        INPUT "Right matrix"; FILER$
580
        INPUT "Output matrix"; FILEO$
590
600
        IF OPT$ = "+" THEN 620
610
        INPUT "a € R
                       in «LEFT*RIGHT"; ALPHA : GOTO 630
        INPUT "\alpha.\beta \in R in \alphaLEFT + \betaRIGHT"; ALPHA, BETA
620
630
        FILE$ = FILEL$ : GOSUB 950
640
        ROW1 = ROW : COL1 = COL : V1 = V
650
        FILE$ = FILER$ : GOSUB 950
660
        ROW2 = ROW : COL2 = COL : V2 = V
670
        IF V1 <> V2 THEN 1730
680
        IF ((OPT$ = "+") AND
        ((ROW1 <> ROW) OR (COL1 <> COL))) THEN 1560
        IF ((OPT = "*") AND (COL1 <> ROW)) THEN 1560
690
700
        OPEN FILEO$ FOR OUTPUT AS #2
        IF OPT$ = "+" THEN GOSUB 2140
710
720
        PRINT #2, ROW1; COL2; V
730
        FOR I1 = 1 TO ROW1
740
          FOR I2 = 1 TO COL2
750
          PRINT "("; I1; ","; I2; ")"
760
            GOSUB 1800
770
            FOR I3 = 1 TO COL1
```

```
780
              T1 = COEFL(I1, I3) : T2 = COEFR(I3, I2)
790
              FOR I4 = 1 TO T1
                 FOR I5 = 1 TO T2
800
                   PROD = ALPHA * A(I1,I3,I4,0) *
810
                                   B(I3, I2, I5,0)
820
                   FOR I6 = 1 TO V
830
                     E(16) = A(11,13,14,16) + B(13,12,15,16)
840
                     NEXT 16
850
                     GOSUB 1150
840
                   NEXT I5
870
                 NEXT 14
880
               NEXT 13
890
            GOSUB 1870
900
            NEXT 12
910
          NEXT I1
920
      · CLOSE #2
        STOP
930
940
        'Subroutine reads a matrix for +/* routines
950
        OPEN FILE$ FOR INPUT AS #1
960
        INPUT #1, ROW, COL, V
970
        IF ((ROW > ROWS) OR (COL > COLS)) THEN 1620
        FOR I = 1 TO ROW
980
990
           FOR J = 1 TO COL
1000
                 INPUT #1, TERMS
1010
                 IF TERMS > TERM THEN 1680
                 IF FILE$ = FILEL$ THEN COEFL(I,J) = TERMS
1020
1030
                 IF FILE$ = FILER$ THEN COEFR(I,J) = TERMS
                 FOR K = 1 TO TERMS
1040
1050
                     FOR L = 0 TO V
1060
                         IF FILE$ = FILER$ THEN 1080
1070
                         INPUT #1, A(I,J,K,L) : GOTO 1090
1080
                         INPUT #1, B(I,J,K,L)
1090
                         NEXT L
                     NEXT K
1100
1110
                 NEXT J
1120
            NEXT I
         CLOSE #1
1130
1140
         RETURN
1150
        'Pointer ordering subroutine
        I = 1 : CNODE = F(0) : BK = 0
1160
1170
        IF CNODE = 0 THEN 1370
1180
        BASE = BK
1190
        IF POT(CNODE) = E(I) THEN 1430
1200
        BK = CNODE
1210
        CNODE = R(CNODE)
1220
        IF CNODE <> 0 THEN 1190
1230
        NODES = NODES + 1
1240
        POT(NODES) = E(I)
1250
        R(BK) = NODES
1260
        BACK(NODES) = BASE
1270
        BK = NODES
        FOR J = I+1 TO V
1280
1290
            NODES = NODES + 1
```

```
1300
           POT(NODES) = E(J)
1310
           BACK(NODES) = BK
1320
           F(BK) = NODES
           BK = NODES
1330
1340
           NEXT J
       M(BK) = M(BK) + PROD
1350
1360
       GOTO 1490
1370
       NODES = NODES + 1
       F(NODES) = 0
1380
1390
       F(BK) = NODES
1400
       BACK(NODES) = BK
1410
       CNODE = NODES
1420
       POT(CNODE) = E(I)
1430
       BASE = CNODE
1440
       I = I + 1
                  'I is the same as the Depth of a node
       IF (I > V) THEN M(CNODE) = M(CNODE) + PROD :
1450
       GOTO 1490
1460
       BK = CNODE
1470
       CNODE = F(CNODE)
1480
       GOTO 1170
1490
       RETURN
1500
        'Error routine for too many polynomial variates
1510
        PRINT "************
1520
        PRINT CHR$(7): "DIM statement allows only ": VAR;
        "variables.";
1530
        PRINT " Modify DIM statement":
1540
        STOP
1550
        'Error routine for ill-dimensioned matrices
1560
        PRINT "*******************************
1570
        PRINT CHR$(7); "Dimensions incompatable"
        PRINT "
                  "; FILEL*; " is "; STR*(ROW1); " X ";
1580
        STR$(COL1)
1590
        PRINT "
                  ": FILER$ : " is ": STR$(ROW): " X ":
        STR$(COL)
1600
1610
        'Error routine for matrices whose dimensions are
         'too large
        PRINT "***********
1620
        PRINT CHR$(7): "Modify the dimension statement in
1630
        the program"
1640
        PRINT; "The dimensions of "; FILE$; "is ";
        STR$ (ROW);
        PRINT " X "; STR$(COL)
1650
1660
1670
        'Error routine for a polynomial having too many
        terms
        1680
        PRINT CHR$(7); "There are "; TERMS; "terms in
1690
1700
        PRINT "polynomial, whereas the DIM statement
        allocates "
1710
        PRINT TERM; " terms in the polynomial.
        DIM stmt."
```

```
1720
         STOP
1730
         PRINT "***************************
1740
         PRINT CHR$(7); "Polynomial variates in matrices
         aren't the same"
1750
1760
         'Error for finding U,M,V,N matrices
        PRINT CHR$(7); "Source matrix "; FILE1$; " doesn't
1770
        have even";
1780
        PRINT " row and column dimensions"
1790
        STOP
1800
        'Subroutine to clear tree building variables
1810
        FOR I = 0 TO NODES
            F(I) = 0 : POT(I) = 0 : R(I) = 0
1820
            BACK(I) = 0 : M(I) = 0
1830
1840
            NEXT I
1850
        NODES = 0
        RETURN
1860
1870
        'Subroutine prints polynomial term
1880
        FOR I = 1 TO NODES 'find # of terms in polynomial
1890
            IF M(I) \iff 0 THEN S = S + 1
1900
1910
            NEXT I
1920
        IF S = 0 THEN 1940
        PRINT #2.5 : GOTO 2000
1930
        PRINT #2,"1";
1940
1950
        FOR I = 0 TO V
                       'handles a zero entry in result
1960
            PRINT #2,0;
1970
            NEXT I
        PRINT #2,""
1980
1990
        GOTO 2130
2000
        FOR I = 1 TO NODES
            IF M(I) = 0 THEN 2120
2010
                                         'coefficient of term
2020
            PRINT #2, M(I);
2030
            PTR = I
2040
            FOR J = V TO 1 STEP -1 'because we are going
                                     from the
2050
                ZZ(J) = POT(PTR)
                                    'top of the tree down
                                     but are
2060
                PTR = BACK(PTR)
                                    'writing the exponents
                                     in the
2070
                NEXT J
                                    'reverse order
2080
            FOR J = 1 TO V
2090
                PRINT #2, ZZ(J);
2100
                NEXT J
            PRINT #2,""
2110
            NEXT I
2120
2130
        RETURN
2140
        'Addition subroutine
2150
        PRINT #2,ROW1; COL1; V1
2160
        FOR I1 = 1 TO ROW1
2170
          FOR I2 = 1 TO COL2
             PRINT "("; I1; ","; I2; ")"
2180
2190
             GOSUB 1800
                                      'clear tree
```

```
2200
        'Send A(I1,I2) up through tree
                FOR I4 = 1 TO COEFL(I1,I2)
2210
2220
                   PROD = ALPHA * A(I1 I2, I4, 0)
2230
                   FOR I6 = 1 TO V
2240
                       E(I6) = A(I1, I2, I4, I6)
2250
                       NEXT 16
2260
                       GOSUB 1150
2270
                   NEXT 14
          'Send B(I1,I2) up through tree
2280
                FOR I4 = 1 TO COEFR(I1,I2)
2290
                   PROD = BETA * B(I1,I2,I4,0)
2300
2310
                   FOR I6 = 1 TO V
                       E(I6) = B(I1, I2, I4, I6)
2320
                       NEXT 16
2330
2340
                       GOSUB 1150
                   NEXT 14
2350
2360
              GOSUB 1870
2370
              NEXT 12
2380
            NEXT I1
2390
          CLOSE #2
2400
          STOP
2410
        'Matrix input subroutine
        INPUT "Matrix name"; FILE$
2420
        INPUT "ROW, COL dimension"; ROW, COL
2430
        INPUT "Number of polynomial variates"; V
2440
        OPEN FILE$ FOR OUTPUT AS #1
2450
2460
        PRINT "Build a square diagonal matrix whose
        diagonal entries ";
2470
        PRINT "are equal Y/N"; : INPUT OPT1$
        IF ((OPT1$ <> "Y") AND (OPT1$ <> "N")) THEN 2460
2480
        PRINT CHR$(7); "*** COEF EXP1 EXP2 ... EXPn ***"
2490
        PRINT " a semicolon; represents end of polynomial"
2500
2510
        PRINT #1, ROW; COL; V
2520
        FOR I = 1 TO ROW
2530
            FOR J = 1 TO COL
                 S = 0 : A = ""
2540
                 IF ((OPT1\$ = "N") OR ((I=1) AND
2550
                 (J=1))) THEN 2610
2560
                 IF I = J THEN S = SS : PRINT #1,S :
                 GOTO 2680
2570
                 PRINT #1.1:
                 FOR K = 0 TO V: PRINT #1,0; : NEXT K
2580
                 PRINT #1.""
2590
2600
                 GOTO 2750
                 PRINT "("; STR$(I); ","; STR$(J); "):
2610
                 WHILE (RIGHT $ (A$,1) <> ";")
2620
2630
                       S = S + 1
                       INPUT A$
2640
                       B$(S) = A$ : BB$(S) = A$
2650
                       WEND
2660
2670
                 PRINT #1,S : SS = S
2680
                 FOR K = 1 TO S-1
                     IF OPT1 = "Y" THEN B $ (K) = BB $ (K)
2690
```

```
PRINT #1,B$(K)
2700
2710
                    NEXT K
                IF OPT1$ = "Y" THEN B$(S) = BB$(S)
2720
2730
                PRINT #1, LEFT$ (B$(S), LEN(B$(S))-1)
2740
                PRINT
2750
                NEXT J
            NEXT I
2760
2770
        CLOSE #1
2780
        GOTO 420
2790
        'subroutine views a matrix file
2800
        INPUT "Screen output Y/N"; OPT1$
        IF ((OPT1$ <> "Y") AND (OPT1$ <> "N")) THEN 2800
2810
        INPUT "Matrix to view"; FILE$
2820
2830
        OPEN FILE$ FOR INPUT AS #1
2840
        INPUT #1, ROW, COL, V
2850
        IF OPT1$ = "Y" THEN 2890
2860
        LPRINT "MATRIX "; FILE$; " IS "; STR$(ROW);
        " X": STR$(COL):
        LPRINT " and has "; STR$(V); " polynomial variates"
2870
        GOTO 2920
2880
        PRINT "MATRIX "; FILE$; " IS "; STR$(ROW); " X";
2890
        STR$(COL);
2900
        PRINT " and has "; STR$(V); " polynomial variates"
2910
        PRINT
2920
        FOR I = 1 TO ROW
2930
            FOR J = 1 TO COL
            IF OPT1$ = "Y" THEN 2970
2940
            LPRINT FILE*; "("; STR*(I); ","; STR*(J); "): "
2950
2960
            GOTO 2980
            PRINT FILE$; "("; STR$(I); ","; STR$(J); "):
2970
            INPUT #1,5
2980
2990
                FOR K = 1 TO S
3000
                    LINE INPUT #1, As
                     IF OPT1$ = "Y" THEN 3030
3010
                     LPRINT A$ : GOTO 3040
3020
3030
                     PRINT A$
3040
                     NEXT K
                IF OPT1$ = "Y" THEN 3080
3050
3060
                LPRINT
3070
                GOTO 3090
3080
                PRINT
3090
                NEXT J
3100
            NEXT I
        CLOSE #1
3110
3120
        GOTO 420
3130
        'Subroutine transposes a matrix
3140
        INPUT "Matrix to transpose"; FILE$
        INPUT "Matrix to store transpose in"; FILEO$
3150
3160
        OPEN FILE$ FOR INPUT AS #1
3170
        INPUT #1, ROW, COL, V
3180
        IF ((ROW > ROWS) OR (COL > COLS)) THEN 1620
        FOR I = 1 TO ROW
3190
3200
            FOR J = 1 TO COL
```

```
3210
                INPUT #1, TERMS
3220
                IF TERMS > TERM THEN 1680
3230
                COEF(I,J) = TERMS
                FOR K = 1 TO TERMS
3240
                     FOR L = 0 TO V
3250
                         INPUT #1, A(I,J,K,L)
3260
3270
                         NEXT L
3280
                    NEXT K
3290
                NEXT J
3300
            NEXT I
         CLOSE #1
3310
3320
         OPEN FILEO$ FOR OUTPUT AS #1
3330
        PRINT #1,COL;ROW;V
3340
        FOR I = 1 TO COL
            FOR J = 1 TO ROW
3350
                PRINT #1, COEF(J,I)
3360
                FOR K = 1 TO COEF(J,I)
3370
                     FOR L = 0 TO V
3380
                         PRINT #1, A(J,I,K,L);
3390
3400
                         NEXT L
                         PRINT #1,""
3410
3420
                     NEXT K
3430
                NEXT J
            NEXT I
3440
         CLOSE #1
3450
        GOTO 420
3460
3470
        'Extracts U,M,V,N matrices
        INPUT "Source matrix of U,M,V,N"; FILE1$
3480
        INPUT "Disk to output to A/B"; D$
3490
        IF ((D$ <> "A") AND (D$ <> "B")) THEN 3490
3500
        D$ = D$ + ":"
3510
        PRINT CHR$(7); "***
                              WARNING: files will be
3520
        output to "; D$;
3530
        PRINT "U,M,V,N ***
        INPUT "OK Y/N"; OPT$
3540
3550
        IF OPT$ <> "Y" THEN STOP
        FILE2* = D* + "U"
3560
        FILE3$ = D$ + "M"
3570
3580
        FILE4$ = D$ + "V"
        FILE5$ = D$ + "N"
3590
        OPEN FILE1$ FOR INPUT AS #1
3600
3610
        OPEN FILE2$ FOR OUTPUT AS #2
        OPEN FILE3$ FOR OUTPUT AS #3
3620
        OPEN FILE4$ FOR OUTPUT AS #4
3630
3640
        OPEN FILES$ FOR OUTPUT AS #5
3650
        INPUT #1,ROW,COL,V
3660
        ROW1 = ROW/2 : COL1 = COL/2
        IF ((ROW1 <> INT(ROW1)) OR
3670
        (COL1 <> INT(COL1))) THEN 1760
        FOR I = 2 TO 5 : PRINT #I, ROW1, COL1, V : NEXT I
3680
3690
        FOR I = 1 TO ROW
3700
            FOR J = 1 TO COL
3710
                 INPUT #1,S
```

```
3720
                FIL = 5
3730
                IF (I <= ROW1) THEN FIL = 2
                IF ((FIL = 2) AND (J > COL1)) THEN FIL = 3
3740
                IF ((FIL = 5) AND (J <= COL1)) THEN FIL = 4
3750
3760
                PRINT #FIL,S
                FOR K = 1 TO S
3770
                    LINE INPUT #1,A$
3780
3790
                    PRINT #FIL, A$
3800
                    NEXT K
3810
                NEXT J
3820
            NEXT I
```

<u>Bibliography</u>

- 1. Ayres, Frank Jr. <u>Schaum's Outline Series Theory and Problems Of Matrices</u>. New York: McGraw Hill Book Company (1962).
- 2. Bellman, R. and R. Vasudevan. "Quasi-linearization and the Matrix Riccati Equation," <u>Journal of Mathematical Analysis and Applications.64</u>: 106-113 (1978).
- 3. Burgess, H.T. "A Practical Method of Determining Elementary Divisors," <u>Annals of Mathematics. second series 18:</u> 4-6 (1916-1917).
- 4. Casti, John L. <u>Dynamical Systems and Their</u>
 <u>Applications</u>. New York: Academic Press (1977).
- 5. Conte, S.D. and Carl de Boor. <u>Elementary Numerical</u>
 <u>Analysis: An Algorithmic Approach</u> (Third Edition).
 New York: McGraw-Hill Book Company (1980).
- One of the sister of the siste
- 7. Fraleigh, John B. <u>A First Course In Abstract Algebra</u> (Second Edition). Reading, Massachusetts: Addison-Wesley Publishing Company (1976).
- 8. Frost, M.G. and C. Storey. "Equivalence of a matrix over R[s,z] with its Smith form," <u>International Journal Of Control. 28</u>: 665-671 (1978).
- 9. Geromel, J.C. "On the Determination of a Diagonal Solution of the Lyapunov Equation," <u>IEEE Transactions</u>
 <u>On Automatic Control. AC-30</u>: 404-406 (April 1985).
- 10. Jensen, Paul A. and J. Wesley Barnes. <u>Network Flow Programming</u>. New York: John Wiley & Sons (1980).
- 11. Jones, John Jr. "Matrix Equations Over Rings,"

 <u>Mathematics and Computers In Simulation, XXV</u>: 489-492

 (1983).
- 12. ----. "Solution Of Certain Matrix Equations,"

 <u>Proceedings of the American Mathematical Society. 31:</u>
 333-339 (February 1972).

13. ----. "On The Matric Equation X2+AX+XB+C=0,"

Proceedings of the American Mathematical Society, 1:
586-589 (1950).

0

- 14. ----. "Computation of the Performance Index of Discrete Linear Time-Invariant Control Systems,"

 <u>Proceedings of the IEEE National Aerospace and Electronics Conference-NAECON</u>: 466-472 (May 1985).
- 15. ----. "Solutions of Algebraic Matrix Equations Related to Optimal Control Theory," <u>Proceedings of a University of Florida International Symposium</u>: 437-440 (1977).
- 16. ---- "Solution of the higher-order Riccati-type matrix equations", <u>International Journal of Systems</u>
 Science, 15 No. 9: 1001-1010 (1984).
- 17. Jones, John Jr et al. "Solutions of Equations A(r)X(r) = b(r)," AIAA (DAY/CIN): 7-3-1--7-3-4 (January 1984).
- 18. Lancaster, Peter. "Explicit Solutions Of Linear Matrix Equations," <u>SIAM Review. 12</u>: 544-549 (October 1970).
- 19. ---- Theory of Matrices. New York: Academic Press, 1969.
- 20. Lee, Bruce E. and Stanislaw H. Zak. "Smith Forms Over R[z₁,z₂]," <u>IEEE Transactions On Automatic Control.</u> AC-28: 115-118 (January 1983).
- 21. Lovass-Nagy, Victor et al. "An Introduction to the Application of the Simplest Matrix-Generalized Inverse in Systems Science," <u>IEEE Transaction on Circuits and Systems</u>, CAS-25: 766-771 (September 1978).
- 22. Macloskie, G. "A General Method of Evaluating Determinants," <u>Annals of Mathematics</u>. 6: 30 (October 1904).
- 23. McClellan, Michael T. "The Exact Solution of Linear Equations With Rational Function Coefficients," ACM Transactions on Mathematical Software. 3: 1-25 (March 1977).
- 24. Moore, E.H. "On the reciprocal of the general algebraic matrix," <u>Bulletin of the American</u>
 <u>Mathematical Society.26</u>: 394-395 (1920).

25. Morris, Gerald L. and Patrick L. Odell. "Common Solutions for n Matrix Equations With Applications,"

Journal of the Association for Computing Machinery, 15: 272-274 (April 1968).

U •

- 26. Nering, Evar D. <u>Linear Algebra and Matrix Theory</u> (second edition). New York: John Wiley & Sons, 1970.
- 27. Pell, A.J. and R.L. Gordon. "The Modified Remainders Obtained in Finding the Highest Common Factor of Two Polynomials," <u>Annals of Mathematics. second series 18:</u> 188-193 (1916-1917).
- 28. Penrose, R. "A generalized inverse for matrices,"

 <u>Proceedings of the Cambridge Philosophical Society. 51:</u>
 406-413 (1955).
- 29. Rao, R. and S.K. Mitra. <u>Generalized Inverses of Matrices and Its Applications</u>. New York: John Wiley & Sons (1971).
- 30. Roth, William E. "The Equations AX YB = C and AX XB = C In Matrices," <u>Proceedings of the American Mathematical Society. 3</u>: 392-396 (1952).
- 31. Sontag, Eduardo D. "On Generalized Inverses of Polynomial and Other Matrices," <u>IEEE Transactions On Automatic Control. AC-25</u>: 514-517 (June 1982).
- 32. Vasudevan, R. "On a Generalization of the Matrix and Scalar Riccati Equation," <u>Journal of Mathematical</u>
 <u>Analysis and Applications.45</u>: 629-638 (1974).
- 33. Wang, Paul S. "Factoring Multivariate Polynomials over Algebraic Number Fields," <u>Mathematics of Computation</u>. 30: 324-336 (April 1976).

VITA

Captain Bruce W. Colletti was born 12 December 1957 in Tucson, Arizona. He graduated from West Phoenix High School in Phoenix, Arizona in May 1976. Attending Arizona State University on a four-year Air Force ROTC Scholarship, he graduated with a Bachelor of Science Degree in Pure and Applied Mathematics in May 1980. Entering active duty one month later, his initial assignment was as an analyst first to the Support Officer Assignments Branch (MPCROS), and then to the Officer Force Analysis Branch (MPCROF), Headquarters Air Force Manpower and Personnel Center, Randolph AFB, Texas. His tour ended in May 1984 upon coming to AFIT for a masters degree in operations research. Decorations include the Air Force Commendation Medal.

Permanent address: 2827 W. Acoma
Phoenix, AZ 85023

	REPORT DOCUM	MENTATION PAGE	E			
Unclassified	1b. RESTRICTIVE MARKINGS					
28 SECURITY CLASSIFICATION AUTHORITY		3. Distribution/Availability of REPORT Approved for public release;				
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		distribution unlimited.				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AFIT/GOR/MA/85D-1		5. MONITORING ORGANIZATION REPORT NUMBER(S)				
6. NAME OF PERFORMING ORGANIZATIO School of Engineering	78. NAME OF MONITORING ORGANIZATION					
Air Force Institute of Tech AFIT/ENG 6c. ADDRESS (City, State and ZIP Code)		7b. ADDRESS (City, State and ZIP Code)				
Wright Patterson AFB, Ohio						
ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER				
8c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUN	IDING NOS.			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT	
11. TITLE (Include Security Classification) See Box 19 (unclassified)						
PERSONAL AUTHOR(S)						
Bruce W. Colletti, CPT. US. 13a TYPE OF REPORT MS Thesis FROM	14. DATE OF REPORT (Yr., Mo., Day) 15. PAGE COUNT December 1985 112					
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES	Continue on reverse if ne	cassam and identic	to by block numb			
FIELD GROUP SUB. GR.	LD GROUP SUB. GR. Riccati equat					
12 01	\longrightarrow equations, mat	ion, Lypanov equation, nonlinear matrix rix equations, polynomial matrices				
19. ABSTRACT (Continue on reverse if necessar Title: "SOLUTIONS OF NONL! Thesis Advisor: John Jones Professor	INEAR MATRIX EQUAT	IONS"	E. WCLAVER for Research and orce Institute of Teat- at-Patterson AFB C	Professional Deve schoology (ATG)	re 🗀	
DISTRIBUTION/AVAILABILITY OF ABST		21. ABSTRACT SECUL		ATION		
224 NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE NU	MBER]	22c OFFICE SYMBOL		
Dr. John Jones Jr		(Include Area Cod. 513 255 3098		AFIT/ENG		

SECURITY CLASSIFICATION OF THIS PAGE

This paper seeks the solutions to a system of equations (equalities) in n variables by expressing the system in matrix algebraic form. Properties of the solutions to the ensuing matrix equation are investigated using similarity transformations. The three types of matrix equations to be studied are the linear equation AX = b, the Lypanov equation AX - XB = C, the second-order Riccati equation XDX + AX + XB + C = 0, and the third-order Riccati equation XAXBX + XCX + DX + XE + F = 0. The entries of all matrices, including the solution X, are restricted to being polynomials in Γ having complex coefficients, where Γ is the n-tuple of indeterminates. That is, all matrices are elements of the ring

 $C_{mXn}(\Gamma)$

for m and n of appropriate size.

Because adding and multiplying matrices (having multivariate polynomial entries) is tedious in practice, an interactive BASIC program is presented in the appendix. This program, which can be used on a personal computer, permits the user to perform operations on matrices having multivariate polynomial entries. Via menu selections, the user may perform

- -weighted addition between two matrices
- -multiplication between two matrices
- -create matrices, with an option of building a diagonal matrix whose diagonal entries are all equal
- -view matrices
- -transpose a matrix
- "xtract special submatrices (U,M,V,N of Chapter IV) from a given matrix

F. N D FILMED) 5-86 DTIC